



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, 5/10, 1/21, C07K 14/705, 14/72, 16/28, A61K 38/17, C12Q 1/68		A2	(11) International Publication Number: WO 00/20590	
			(43) International Publication Date: 13 April 2000 (13.04.00)	
(21) International Application Number: PCT/US99/23317		<p>Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). AU-YOUNG, Janice [US/US]; 1419 Kains Avenue, Berkeley, CA 94702 (US). REDDY, Roopa [IN/US]; 1233 W. McKinley Drive, Sunnyvale, CA 94086 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue #30, Mountain View, CA 94040 (US). GUEGLER, Karl, J. [CH/US]; 1048 Oakland Avenue, Menlo Park, CA 94025 (US). GORGONE, Gina, A. [US/US]; 1253 Pinecrest Drive, Boulder Creek, CA 95006 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US).</p> <p>(74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p>		
(22) International Filing Date: 6 October 1999 (06.10.99)				
(30) Priority Data:				
09/167,219	6 October 1998 (06.10.98)			US
60/172,211	6 October 1998 (06.10.98)			US
60/133,585	11 May 1999 (11.05.99)	US		
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications				
US	09/167,219 (CIP)			
Filed on	6 October 1998 (06.10.98)			
US	60/172,211 (CIP)			
Filed on	6 October 1998 (06.10.98)			
US	60/133,585 (CIP)			
Filed on	11 May 1999 (11.05.99)			
(71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US).				
(72) Inventors; and				
(75) Inventors/Applicants (for US only): TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). YUE,				

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: G-PROTEIN COUPLED RECEPTOR PROTEINS

```

1  M-----T P Q S L L Q T T L F L L S L-----L F L V 2763296
1  M-----T P Q S L L Q T T L F L L S L-----L F L V 1595762
1  M V F S V R Q C G H V G R T E E V L L T F K I F L V I I C L H V V L V GI 2117161

21 QGAHGRGRHREDPR-----FCSQRNQTHRSS 2763296
21 QGAHGRGRHREDPR-----FCSQRNQTHRSS 1595762
36 T S L E E D T D N S S L S P P P A K L S V V S F A P S S N E V E T T S GI 2117161

46 L H Y-----2763296
46 L H Y K P T P D L R I S I E N S E E A L T V H A P P P A A H P A S R-----1595762
71 L N - D V T L S L L P S N E T E K T K I T I V R T F N A S G V K P Q R GI 2117161

49 -----2763296
80 -----C L Y W N R H-----1595762
105 N I C N L S S I C N D S A F P R G E I M F Q Y D K E S T V P Q N Q H I GI 2117161

49 -----2763296
98 -----A G R L H L L Y G K R D-----F L L-----1595762
140 T N G T L T G V L S L S E L K R S E L N K T L Q T L S E T Y P I M C A GI 2117161

49 -----2763296
113 -S D K A S S L L C-----1595762
175 T A E A Q S T L N C T P T I K L N N T M N A C A I A A L E R V K I R GI 2117161

```

(57) Abstract

The invention provides human G-protein coupled receptor proteins (GCRP) and polynucleotides which identify and encode GCRP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of GCRP.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

G-PROTEIN COUPLED RECEPTOR PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of G-protein coupled receptor proteins and to the use of these sequences in the diagnosis, treatment, and prevention of disorders of the nervous system, autoimmune/inflammatory disorders, and cell proliferative disorders including cancer.

BACKGROUND OF THE INVENTION

Regulation of cell proliferation, differentiation, and migration is important for the formation and function of tissues. Regulatory proteins such as growth factors control these cellular processes and act as mediators in cell-cell signaling pathways. Growth factors are secreted proteins that bind to specific cell-surface receptors on target cells. The bound receptors trigger intracellular signal transduction pathways which activate various downstream effectors that regulate gene expression, cell division, cell differentiation, cell motility, and other cellular processes. Some of the receptors involved in signal transduction by growth factors belong to the large superfamily of G-protein coupled receptors (GPCRs).

GPCRs are integral membrane proteins characterized by the presence of seven hydrophobic transmembrane domains which together form a bundle of antiparallel alpha (α) helices. These proteins range in size from under 400 to over 1000 amino acids (Strosberg, A.D. (1991) Eur. J. Biochem. 196:1-10; Coughlin, S.R. (1994) Curr. Opin. Cell Biol. 6:191-197). The amino-terminus of a GPCR is extracellular, is of variable length, and is often glycosylated. The carboxy-terminus is cytoplasmic and generally phosphorylated. Extracellular loops of GPCRs alternate with intracellular loops and link the transmembrane domains. Cysteine disulfide bridges linking the second and third extracellular loops may interact with agonists and antagonists. The most conserved domains of GPCRs are the transmembrane domains and the first two cytoplasmic loops. The transmembrane domains account for structural and functional features of the receptor. In most cases, the bundle of α helices forms a ligand-binding pocket. The extracellular N-terminal segment, or one or more of the three extracellular loops, may also participate in ligand binding. Ligand binding activates the receptor by inducing a conformational change in intracellular portions of the receptor. In turn, the large, third intracellular loop of the activated receptor interacts with an intracellular heterotrimeric guanine nucleotide binding (G) protein complex which mediates further intracellular signaling activities, including the activation of second messengers such as cyclic AMP (cAMP), phospholipase C, inositol triphosphate, or ion channel proteins. (See, e.g., Watson, S. and S. Arkinstall (1994) The G-protein Linked Receptor Facts Book, Academic Press, San Diego CA, pp. 2-6; Bolander, F.F. (1994) Molecular Endocrinology, Academic Press, San Diego CA, pp. 162-176; Baldwin, J.M. (1994) Curr. Opin. Cell Biol. 6:180-190.)

GPCRs include receptors for sensory signal mediators (e.g., light and olfactory stimulatory

molecules); adenosine, bombesin, bradykinin, endothelin, γ -aminobutyric acid (GABA), hepatocyte growth factor, melanocortins, neuropeptide Y, opioid peptides, opsins, somatostatin, tachykinins, vasoactive intestinal polypeptide family, and vasopressin; biogenic amines (e.g., dopamine, epinephrine and norepinephrine, histamine, glutamate (metabotropic effect), acetylcholine (muscarinic effect), and serotonin); chemokines; lipid mediators of inflammation (e.g., prostaglandins and prostanoids, platelet activating factor, and leukotrienes); and peptide hormones (e.g., calcitonin, C5a anaphylatoxin, follicle-stimulating hormone (FSH), gonadotropic-releasing hormone (GnRH), neurokinin, and thyrotropin-releasing hormone (TRH), and oxytocin). GPCRs which act as receptors for stimuli that have yet to be identified are known as orphan receptors.

GPCRs are implicated in inflammation and the immune response, and include the EGF module-containing, mucin-like hormone receptor (Emr1) and CD97 β receptor proteins. These receptors contain between three and seven potential calcium-binding EGF-like motifs (Baud, V. et al. (1995) *Genomics* 26:334-344; Gray, J.X. et al. (1996) *J. Immunol.* 157:5438-5447). These GPCRs are members of the recently characterized EGF-TM7 receptors family. In addition, post-translational modification of aspartic acid or asparagine to form erythro- β -hydroxyaspartic acid or erythro- β -hydroxyasparagine has been identified in a number of proteins with domains homologous to EGF. The consensus pattern is located in the N-terminus of the EGF-like domain. Examples of such proteins are blood coagulation factors VII, IX, and X; proteins C, S, and Z; the LDL receptor; and thrombomodulin.

One large subfamily of GPCRs are the olfactory receptors. These receptors share the seven hydrophobic transmembrane domains of other GPCRs and function by registering G protein-mediated transduction of odorant signals. Numerous distinct olfactory receptors are required to distinguish different odors. Each olfactory sensory neuron expresses only one type of olfactory receptor, and distinct spatial zones of neurons expressing distinct receptors are found in nasal passages. One olfactory receptor, the RA1c receptor which was isolated from a rat brain library, has been shown to be limited in expression to very distinct regions of the brain and a defined zone of the olfactory epithelium (Raming, K. et al. (1998) *Receptors Channels* 6:141-151). In another example, three rat genes encoding olfactory-like receptors having typical GPCR characteristics showed expression patterns exclusively in taste, olfactory, and male reproductive tissue (Thomas, M.B. et al. (1996) *Gene* 178:1-5).

Another group of GPCRs are the mas oncogene-related proteins. Like the mas oncogenes themselves, some of these mas-like receptors are implicated in intracellular angiotensin II actions. Angiotensin II, an octapeptide hormone, mediates vasoconstriction and aldosterone secretion through angiotensin II receptor molecules found on smooth vascular muscle and the adrenal glands, respectively. A cloned human mas-related gene (mrg) mRNA, when injected into *Xenopus* oocytes, produces an increase in the response to angiotensin peptides. Mrg has been shown to directly affect signaling pathways associated with the angiotensin II receptor, and, accordingly, affects the processes of

vasoconstriction and aldosterone secretion (Monnot, C. et al. (1991) Mol. Endocrinol. 5:1477-1487).

GPCR mutations, which may cause loss of function or constitutive activation, have been associated with numerous human diseases (Coughlin, *supra*). For instance, retinitis pigmentosa may arise from mutations in the rhodopsin gene. Rhodopsin is the retinal photoreceptor which is located within the discs of the eye rod cell. Parma, J. et al. (1993, Nature 365:649-651) reported that somatic activating mutations in the thyrotropin receptor cause hyperfunctioning thyroid adenomas and suggested that certain GPCRs susceptible to constitutive activation may behave as protooncogenes.

The discovery of new G-protein coupled receptor proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of disorders of the nervous system, autoimmune/inflammatory disorders, and cell proliferative disorders including cancer.

SUMMARY OF THE INVENTION

The invention features substantially purified polypeptides, G-protein coupled receptor proteins, referred to collectively as "GCRP" and individually as "GCRP-1," "GCRP-2," "GCRP-3," "GCRP-4," "GCRP-5," "GCRP-6," "GCRP-7," and "GCRP-8." In one aspect, the invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof.

The invention further provides a substantially purified variant having at least 90% amino acid identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-8, and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof.

The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates

with the presence of a polynucleotide in the sample. In one aspect, the method further comprises amplifying the polynucleotide prior to hybridization.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9-16, and fragments thereof.

10 The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof. In another aspect, the expression vector is contained within a host cell.

15 The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing at least a fragment of a polynucleotide under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

20 The invention further includes a purified antibody which binds to a polypeptide selected from the group consisting of SEQ ID NO:1-8, and fragments thereof. The invention also provides a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a disorder associated with decreased expression or activity of GCRP, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

25 The invention also provides a method for treating or preventing a disorder associated with increased expression or activity of GCRP, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-8, and fragments thereof.

BRIEF DESCRIPTION OF THE FIGURES AND TABLES

35 Figures 1A, 1B, 1C, 1D, and 1E show the amino acid sequence alignment between GCRP-2

(2763296; SEQ ID NO:2), GCRP-1 (1595762; SEQ ID NO:1), and human epididymis-specific seven transmembrane receptor HE6 (GI 2117161; SEQ ID NO:17), produced using the multisequence alignment program of LASERGENE software (DNASTAR, Madison WI).

Figures 2A and 2B show the amino acid sequence alignment between GCRP-3 (3367641; SEQ ID NO:3) and human CD97 (GI 3342740; SEQ ID NO:18), produced using the multisequence alignment program of LASERGENE software.

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding GCRP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods and algorithms used for identification of GCRP.

Table 3 shows useful fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding GCRP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze GCRP, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in

connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

“GCRP” refers to the amino acid sequences of substantially purified GCRP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and preferably the human species, from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term “agonist” refers to a molecule which, when bound to GCRP, increases or prolongs the duration of the effect of GCRP. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of GCRP.

An “allelic variant” is an alternative form of the gene encoding GCRP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding GCRP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide the same as GCRP or a polypeptide with at least one functional characteristic of GCRP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding GCRP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding GCRP. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent GCRP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of GCRP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. In this context, “fragments,” “immunogenic fragments,” or “antigenic fragments” refer to fragments of

GCRP which are preferably at least 5 to about 15 amino acids in length, most preferably at least 14 amino acids, and which retain some biological activity or immunological activity of GCRP. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which, when bound to GCRP, decreases the amount or the duration of the effect of the biological or immunological activity of GCRP. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies, or any other molecules which decrease the effect of GCRP.

The term "antibody" refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies that bind GCRP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that fragment of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (given regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" can refer to the antisense strand, and the designation "positive" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic GCRP, or of any oligopeptide thereof, to induce a specific

immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such
5 that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

10 A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding GCRP or fragments of GCRP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated
15 with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3'
20 direction, and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte clone using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

The term "correlates with expression of a polynucleotide" indicates that the detection of the
25 presence of nucleic acids, the same or related to a nucleic acid sequence encoding GCRP, by northern analysis is indicative of the presence of nucleic acids encoding GCRP in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding GCRP.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

30 The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at
35 least one biological or immunological function of the polypeptide from which it was derived.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern
5 or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as
10 reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

15 The phrases "percent identity" and "% identity" refer to the percentage of sequence similarity found in a comparison of two or more amino acid or nucleic acid sequences. Percent identity can be determined electronically, e.g., by using the MEGALIGN program (DNASTAR, Madison WI) which creates alignments between two or more sequences according to methods selected by the user, e.g., the clustal method. (See, e.g., Higgins, D.G. and P.M. Sharp (1988) *Gene* 73:237-244.) Parameters for each
20 method may be the default parameters provided by MEGALIGN or may be specified by the user. The clustal algorithm groups sequences into clusters by examining the distances between all pairs. The clusters are aligned pairwise and then in groups. The percentage similarity between two amino acid sequences, e.g., sequence A and sequence B, is calculated by dividing the length of sequence A, minus the number of gap residues in sequence A, minus the number of gap residues in sequence B, into the sum
25 of the residue matches between sequence A and sequence B, times one hundred. Gaps of low or of no similarity between the two amino acid sequences are not included in determining percentage similarity. Percent identity between nucleic acid sequences can also be counted or calculated by other methods known in the art, e.g., the Jotun Hein method. (See, e.g., Hein, J. (1990) *Methods Enzymol.* 183:626-645.) Identity between sequences can also be determined by other methods known in the art, e.g., by
30 varying hybridization conditions.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence
35 in the non-antigen binding regions has been altered so that the antibody more closely resembles a human

antibody, and still retains its original binding ability.

"Hybridization" refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

10 The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, to the sequence found in the naturally occurring molecule.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

20 The term "modulate" refers to a change in the activity of GCRP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of GCRP.

The phrases "nucleic acid" or "nucleic acid sequence," as used herein, refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material. In this context, "fragments" refers to those nucleic acid sequences which comprise a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:9-16, for example, as distinct from any other sequence in the same genome. For example, a fragment of SEQ ID NO:9-16 is useful in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:9-16 from related polynucleotide sequences. A fragment of SEQ ID NO:9-16 is at least about 15-20 nucleotides in length. The precise length of the fragment of SEQ ID NO:9-16 and the region of SEQ ID NO:9-16 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment. In some cases, a fragment, when translated, would produce polypeptides retaining some functional characteristic, e.g., antigenicity, or structural domain

characteristic, e.g., ATP-binding site, of the full-length polypeptide.

The terms "operably associated" and "operably linked" refer to functionally related nucleic acid sequences. A promoter is operably associated or operably linked with a coding sequence if the promoter controls the translation of the encoded polypeptide. While operably associated or operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements, e.g., repressor genes, are not contiguously linked to the sequence encoding the polypeptide but still bind to operator sequences that control expression of the polypeptide.

The term "oligonucleotide" refers to a nucleic acid sequence of at least about 6 nucleotides to 60 nucleotides, preferably about 15 to 30 nucleotides, and most preferably about 20 to 25 nucleotides, which can be used in PCR amplification or in a hybridization assay or microarray. "Oligonucleotide" is substantially equivalent to the terms "amplimer," "primer," "oligomer," and "probe," as these terms are commonly defined in the art.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding GCRP, or fragments thereof, or GCRP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "stringent conditions" refers to conditions which permit hybridization between polynucleotides and the claimed polynucleotides. Stringent conditions can be defined by salt concentration, the concentration of organic solvent, e.g., formamide, temperature, and other conditions well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally

associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of GCRP polypeptides refers to an amino acid sequence that is altered by one or more amino acid residues. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have "nonconservative" changes (e.g., replacement of glycine with tryptophan). Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, LASERGENE software (DNASTAR).

The term "variant," when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to GCRP. This definition may also include, for example, "allelic" (as defined above), "splice," "species," or "polymorphic" variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity

for a disease state.

THE INVENTION

The invention is based on the discovery of new human G-protein coupled receptor proteins (GCRP), the polynucleotides encoding GCRP, and the use of these compositions for the diagnosis, treatment, or prevention of disorders of the nervous system, autoimmune/inflammatory disorders, and cell proliferative disorders including cancer.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding GCRP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each GCRP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The clones in column 5 were used to assemble the consensus nucleotide sequence of each GCRP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs, as well as GenBank homologs identified by BLAST analysis; column 6 shows the identity of each polypeptide; and column 7 shows analytical methods used to characterize each polypeptide through sequence homology and protein motifs.

As shown in Figures 1A, 1B, 1C, 1D, and 1E, GCRP-1 has chemical and structural similarity with human epididymis-specific seven transmembrane receptor HE6 (GI 2117161; SEQ ID NO:17). In particular, GCRP-1 and human HE6 share 19% identity, four potential N-glycosylation sites, and one potential protein kinase C phosphorylation site. GCRP-1 and GCRP-2 have chemical and structural homology and are considered to be splice variants. Therefore, GCRP-2 also has chemical and structural similarity with human epididymis-specific seven transmembrane receptor HE6 (GI 2117161; SEQ ID NO:17). As shown in Figures 2A and 2B, GCRP-3 has chemical and structural similarity with human CD97 (GI 3342740; SEQ ID NO:18). In particular, GCRP-3 and human CD97 share 44% identity, one potential N-glycosylation site, one potential protein kinase C phosphorylation site, and one potential aspartic acid/asparagine hydroxylation site.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding GCRP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:9-16, and to

distinguish between SEQ ID NO:9-16 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express GCRP as a fraction of total tissues expressing GCRP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing GCRP as a fraction of total tissues expressing GCRP. Column 5 lists the vectors used to subclone each cDNA library. Northern analysis shows the expression of SEQ ID NO:9 in various libraries, at least 55% of which are associated with cancer and cell proliferation, and at least 30% of which are associated with the immune response. Northern analysis shows the expression of SEQ ID NO:11 in various libraries, at least 50% of which are associated with cancer and at least 50% of which are associated with the immune response. Of particular note is the expression of SEQ ID NO:9 in brain, thyroid, epididymis, prostate, breast, uterus, kidney, lung, gut, and bladder; the expression of SEQ ID NO:10 in brain, kidney, epididymis, prostate, breast, and bladder; the expression of SEQ ID NO:11 in tumor, immunological, and inflammatory tissues; the expression of SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:16 in cancerous and proliferating tissues; the expression of SEQ ID NO:12 in brain tumor; the expression of SEQ ID NO:13 in kidney tumor and prostate tumor; and the expression of SEQ ID NO:14 in tongue tumor and prostate.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding GCRP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses GCRP variants. A preferred GCRP variant is one which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% amino acid sequence identity to the GCRP amino acid sequence, and which contains at least one functional or structural characteristic of GCRP.

The invention also encompasses polynucleotides which encode GCRP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:9-16, which encodes GCRP.

The invention also encompasses a variant of a polynucleotide sequence encoding GCRP. In particular, such a variant polynucleotide sequence will have at least about 70%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding GCRP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:9-16 which has at least about 70%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:9-16. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of GCRP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding GCRP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made
5 by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring GCRP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode GCRP and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring GCRP under appropriately selected
10 conditions of stringency, it may be advantageous to produce nucleotide sequences encoding GCRP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding GCRP
15 and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode GCRP and GCRP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic
20 sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding GCRP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:9-16, and
25 fragments, thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) Methods Enzymol. 152:399-407; Kimmel, A.R. (1987) Methods Enzymol. 152:507-511.) For example, stringent salt concentration will ordinarily be less than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and most preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the
30 absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide, and most preferably at least about 50% formamide. Stringent temperature conditions will ordinarily include temperatures of at least about 30°C, more preferably of at least about 37°C, and most preferably of at least about 42°C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion
35 or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are

accomplished by combining these various conditions as needed. In a preferred embodiment, hybridization will occur at 30°C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37°C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 µg/ml denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will occur at 42°C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50 % formamide, and 200 µg/ml ssDNA. Useful variations on these conditions will be readily apparent to those skilled in the art.

The washing steps which follow hybridization can also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include temperature of at least about 25°C, more preferably of at least about 42°C, and most preferably of at least about 68°C. In a preferred embodiment, wash steps will occur at 25°C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 42°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a most preferred embodiment, wash steps will occur at 68°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. Additional variations on these conditions will be readily apparent to those skilled in the art.

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA), and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using the ABI 373 or 377 DNA sequencing systems (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding GCRP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site

PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and
5 surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other
10 methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis
15 software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include
20 sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary
25 sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled.
30 Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode GCRP may be cloned in recombinant DNA molecules that direct expression of GCRP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of
35 the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent

amino acid sequence may be produced and used to express GCRP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter GCRP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling
5 by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding GCRP may be synthesized, in whole or in part,
10 using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, GCRP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-
15 Elmer). Additionally, the amino acid sequence of GCRP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The
20 composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active GCRP, the nucleotide sequences encoding GCRP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a
25 suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding GCRP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding GCRP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding
30 GCRP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The
35 efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host

cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding GCRP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding GCRP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding GCRP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding GCRP can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding GCRP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When large quantities of GCRP are needed, e.g. for the production of antibodies, vectors which direct high level expression of GCRP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of GCRP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184.)

Plant systems may also be used for expression of GCRP. Transcription of sequences encoding GCRP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311).

Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g.,

5 The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding GCRP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader

10 sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses GCRP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

15 Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of

20 GCRP in cell lines is preferred. For example, sequences encoding GCRP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a

25 selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase

30 genes, for use in *tk* or *ap^r* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci.

35 USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable

genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding GCRP is inserted within a marker gene sequence, transformed cells containing sequences encoding GCRP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GCRP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding GCRP and that express GCRP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of GCRP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on GCRP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding GCRP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding GCRP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available

kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

5 Host cells transformed with nucleotide sequences encoding GCRP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode GCRP may be designed to contain signal sequences which direct secretion
10 of GCRP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may
15 also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

20 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding GCRP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric GCRP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of GCRP activity. Heterologous protein and peptide moieties
25 may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-*
30 *myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the GCRP encoding sequence and the heterologous protein sequence, so that GCRP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are
35 discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to

facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled GCRP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract systems (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, preferably ³⁵S-methionine.

Fragments of GCRP may be produced not only by recombinant production, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of GCRP may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of GCRP and G-protein coupled receptor proteins. In addition, the expression of GCRP is closely associated with cell proliferation, cancer, and autoimmune disease/inflammation. Therefore, GCRP appears to play a role in disorders of the nervous system, autoimmune/inflammatory disorders, and cell proliferative disorders including cancer. In the treatment of disorders associated with increased GCRP expression or activity, it is desirable to decrease the expression or activity of GCRP. In the treatment of disorders associated with decreased GCRP expression or activity, it is desirable to increase the expression or activity of GCRP.

Therefore, in one embodiment, GCRP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCRP. Examples of such disorders include, but are not limited to, a nervous disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease; prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome; fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorder of the central nervous system, cerebral palsy, a neuroskeletal disorder, an autonomic nervous system disorder, a cranial nerve disorder, a spinal cord disease, muscular dystrophy and other neuromuscular disorders, a peripheral nervous system disorder, dermatomyositis and polymyositis; inherited, metabolic, endocrine, and toxic

myopathy; myasthenia gravis, periodic paralysis; a mental disorder including mood, anxiety, and schizophrenic disorders; seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.

In another embodiment, a vector capable of expressing GCRP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCRP including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified GCRP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCRP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GCRP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GCRP including, but not limited to, those listed above.

In a further embodiment, an antagonist of GCRP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GCRP. Examples of such disorders

include, but are not limited to, disorders of the nervous system, autoimmune/inflammatory disorders, and cell proliferative disorders. In one aspect, an antibody which specifically binds GCRP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express GCRP.

5 In an additional embodiment, a vector expressing the complement of the polynucleotide encoding GCRP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GCRP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate
10 therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

15 An antagonist of GCRP may be produced using methods which are generally known in the art. In particular, purified GCRP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GCRP. Antibodies to GCRP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab
20 expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with GCRP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase
25 immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to GCRP
30 have an amino acid sequence consisting of at least about 5 amino acids, and, more preferably, of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of GCRP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

35 Monoclonal antibodies to GCRP may be prepared using any technique which provides for the

production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce GCRP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for GCRP may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between GCRP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering GCRP epitopes is preferred, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for GCRP. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of GCRP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined

for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple GCRP epitopes, represents the average affinity, or avidity, of the antibodies for GCRP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular GCRP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12}

- 5 L/mole are preferred for use in immunoassays in which the GCRP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of GCRP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to
10 Monoclonal Antibodies, John Wiley & Sons, New York NY).

- The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is preferred for use in procedures requiring precipitation of GCRP-antibody
15 complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

- In another embodiment of the invention, the polynucleotides encoding GCRP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the
20 polynucleotide encoding GCRP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding GCRP. Thus, complementary molecules or fragments may be used to modulate GCRP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along
25 the coding or control regions of sequences encoding GCRP.

- Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding GCRP. (See,
30 e.g., Sambrook, supra; Ausubel, 1995, supra.)

- Genes encoding GCRP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding GCRP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled
35 by endogenous nucleases. Transient expression may last for a month or more with a non-replicating

vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding GCRP. Oligonucleotides derived from the transcription initiation site, e.g.,
5 between about positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and
10 Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme
15 molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding GCRP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU,
20 and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

25 Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding GCRP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable
30 RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within
35 the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in

all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of GCRP, antibodies to GCRP, and mimetics, agonists, antagonists, or inhibitors of GCRP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain

tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms.

In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of GCRP, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

10 For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

15 A therapeutically effective dose refers to that amount of active ingredient, for example GCRP or fragments thereof, antibodies of GCRP, and agonists, antagonists or inhibitors of GCRP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the
20 population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this
25 range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of
30 the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μ g to 100,000 μ g, up to a total dose of about 1
35 gram, depending upon the route of administration. Guidance as to particular dosages and methods of

delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

5 In another embodiment, antibodies which specifically bind GCRP may be used for the diagnosis of disorders characterized by expression of GCRP, or in assays to monitor patients being treated with GCRP or agonists, antagonists, or inhibitors of GCRP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for GCRP include methods which utilize the antibody and a label to detect GCRP in human body fluids or in extracts of cells
10 or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring GCRP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of GCRP expression. Normal or
15 standard values for GCRP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to GCRP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, preferably by photometric means. Quantities of GCRP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject
20 values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding GCRP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and
quantitate gene expression in biopsied tissues in which expression of GCRP may be correlated with
25 disease. The diagnostic assay may be used to determine absence, presence, and excess expression of GCRP, and to monitor regulation of GCRP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding GCRP or closely related molecules may be used to identify nucleic acid sequences which encode GCRP. The specificity of the probe, whether it is made
30 from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine whether the probe identifies only naturally occurring sequences encoding GCRP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably have at
35 least 50% sequence identity to any of the GCRP encoding sequences. The hybridization probes of the

subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:9-16 or from genomic sequences including promoters, enhancers, and introns of the GCRP gene.

Means for producing specific hybridization probes for DNAs encoding GCRP include the cloning of polynucleotide sequences encoding GCRP or GCRP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

- Polynucleotide sequences encoding GCRP may be used for the diagnosis of disorders associated with expression of GCRP. Examples of such disorders include, but are not limited to, a nervous disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease; prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome; fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorder of the central nervous system, cerebral palsy, a neuroskeletal disorder, an autonomic nervous system disorder, a cranial nerve disorder, a spinal cord disease, muscular dystrophy and other neuromuscular disorders, a peripheral nervous system disorder, dermatomyositis and polymyositis; inherited, metabolic, endocrine, and toxic myopathy; myasthenia gravis, periodic paralysis; a mental disorder including mood, anxiety, and schizophrenic disorders; seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis,

anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma. The polynucleotide sequences encoding GCRP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered GCRP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GCRP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding GCRP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding GCRP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of GCRP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GCRP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization

assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding GCRP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding GCRP, or a fragment of a polynucleotide complementary to the polynucleotide encoding GCRP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of GCRP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding GCRP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The

sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) site. Correlation between the location of the gene encoding GCRP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, GCRP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between GCRP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with GCRP, or fragments thereof, and washed. Bound GCRP is then detected by methods well known in the art. Purified GCRP can also be coated directly onto

plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GCRP specifically compete with a test compound for binding GCRP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with GCRP.

In additional embodiments, the nucleotide sequences which encode GCRP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. [Attorney Docket No. PF-0612 P, filed October 6, 1998] and U.S. Ser. No. 60/133,585, are hereby expressly incorporated by reference.

20 **EXAMPLES**

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP

vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSORT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA).

10 Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8, QIAWELL 8 Plus, QIAWELL 8 Ultra plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

20 Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing systems (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software;

35

or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:9-16. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene

and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

10

100

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding GCRP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation/trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of GCRP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:9-16 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was

performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

- 10 The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of
- 15 DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

- The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun
- 20 sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media,
- 25 individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

- The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated
- 30 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulphoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction
- 35 kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:9-16 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

VI. Labeling and Use of Individual Hybridization Probes

- 5 Hybridization probes derived from SEQ ID NO:9-16 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine
- 10 triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont
- 15 NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate.

- 20 Hybridization patterns are visualized using autoradiography and compared.

VII. Microarrays

- A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, supra.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or
- 25 mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.
- 30 Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a
- 35 glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and

chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

- 5 Sequences complementary to the GCRP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring GCRP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of GCRP. To inhibit transcription, a
10 complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the GCRP-encoding transcript.

IX. Expression of GCRP

- Expression and purification of GCRP is achieved using bacterial or virus-based expression
15 systems. For expression of GCRP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria
20 express GCRP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of GCRP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding GCRP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates.
25 Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945.)

- 30 In most expression systems, GCRP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech).
35 Following purification, the GST moiety can be proteolytically cleaved from GCRP at specifically

engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16).

5 Purified GCRP obtained by these methods can be used directly in the following activity assay.

X. Demonstration of GCRP Activity

The activity of GCRP may be measured using an assay based upon the property of some GPCRs to support proliferation in vitro of fibroblasts and tumor cells under serum-free conditions (Chiquet-Ehrismann, R. et al. (1986) Cell 47:131-139). Wells in 96 well cluster plates (Falcon, Fisher Scientific,
10 Santa Clara CA) are coated with GCRP by incubation with solutions at 50-100 µg/ml for 15 min at ambient temperature. The coating solution is aspirated, and the wells washed with Dulbecco's medium before cells are plated. Rat fibroblast cultures or rat mammary tumor cells are prepared as described and plated at a density of 10^4 - 10^5 cells/ml in Dulbecco's medium supplemented with 10% fetal calf serum (FCS).
15 After three days the media are removed, and the cells washed three times with phosphate-buffered saline (PBS) before the addition of serum-free Dulbecco's medium containing 0.25 mg/ml bovine serum albumin (BSA, Fraction V, Sigma Chemical, St. Louis, MO). After 2 days the medium is aspirated, and 100 µl of [3H]thymidine (NEN) at 2 µCi/ml in fresh Dulbecco's medium containing 0.25 mg/ml BSA added. Parallel plates are fixed and stained to determine cell numbers. After 16 hr, the
20 medium is aspirated, the cell layer washed with PBS, and the 10% trichloroacetic acid-precipitable counts in the cell layer determined by liquid scintillation counting of radioisotope (normalized to relative cell numbers; Chiquet-Ehrismann, R. et al. (1986) supra). The rates of cell proliferation and [3H]thymidine uptake are proportional to the levels of GCRP in the sample.

In the alternative, the assay for GCRP activity is based upon the property of CD97/Emr1 GPCR
25 family proteins to modulate G protein-activated second messenger signal transduction pathways (e.g., cAMP; Gaudin, P. et al. (1998) J. Biol. Chem. 273:4990-4996). A plasmid encoding full length GCRP is transfected into a mammalian cell line (e.g., COS-7 or Chinese hamster ovary (CHO-K1) cell lines) using methods well-known in the art. Transfected cells are grown in 12-well trays in culture medium containing 2% FCS for 48 hours, the culture medium is discarded, then the attached cells are gently
30 washed with PBS. The cells are then incubated in culture medium with 10% FCS or 2% FCS for 30 minutes, then the medium is removed and cells lysed by treatment with 1 M perchloric acid. The cAMP levels in the lysate are measured by radioimmunoassay using methods well-known in the art. Changes in the levels of cAMP in the lysate from 10% FCS-treated cells compared with those in 2% FCS-treated cells are proportional to the amount of GCRP present in the transfected cells.

35 XI. Functional Assays

GCRP function is assessed by expressing the sequences encoding GCRP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, preferably of endothelial or hematopoietic origin, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of GCRP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding GCRP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding GCRP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of GCRP Specific Antibodies

GCRP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the GCRP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for

selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

10 XIII. Purification of Naturally Occurring GCRP Using Specific Antibodies

Naturally occurring or recombinant GCRP is substantially purified by immunoaffinity chromatography using antibodies specific for GCRP. An immunoaffinity column is constructed by covalently coupling anti-GCRP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed
15 according to the manufacturer's instructions.

Media containing GCRP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of GCRP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/GCRP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and
20 GCRP is collected.

XIV. Identification of Molecules Which Interact with GCRP

GCRP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton et al. (1973) Biochem. J. 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled GCRP, washed, and any wells with labeled
25 GCRP complex are assayed. Data obtained using different concentrations of GCRP are used to calculate values for the number, affinity, and association of GCRP with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.

30 Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	9	1595762	BRAINOT14	1595762H1 (BRAINOT14), 1546740R6 (PROSTUT04), 1597820F6 (BRAINOT14), 2716791F3, (THYRNOT09), 1705908F6 (DUODNOT02), 1753259F6 (LIVRTUT01), 1453415F1 (PENITUT01), 1314567F1 (BLADTUT02), 115604F1 (KIDNNOT01), SBIA04259D1, SBIA03661D1, SBIA05044D1
2	10	2763296	BRSTNOT12	2763296H1 (BRSTNOT12), 3486025F6 (KIDNNOT31), SBDA07002F3
3	11	3367641	CONNTUT04	340333R6 (NEUTFMT01), 3367641F6 (CONNTUT04), 3367641H1 (CONNTUT04), 3367641T6 (CONNTUT04)
4	12	866390	BRAITUT03	866390H1 (BRAITUT03), 866390T1 (BRAITUT03), 2112194R6 (BRAITUT03)
5	13	998550	KIDNTUT01	998550H1 (KIDNTUT01), 998550X312D1 (KIDNTUT01), SXCA00752V1, SCDA06667V1, SXCA00292V1
6	14	1547873	PROSNOT06	980190R6 (TONGTUT01), 980190T1 (TONGTUT01), 1547873H1 (PROSNOT06), SBHA00047F1, SBHA00670F1
7	15	2048484	LIVRFET02	1872334F6 (LEUKNOT02), 1872334T3 (LEUKNOT02), 2048484H1 (LIVRFET02), SBGA01350F1, SBGA04183F1
8	16	5554170	TONSDIT01	1433664H1 (BEPINON01), 1522189F6 (BLADTUT04), 1542382R1 (SINTTUT01), 1662775T6 (BRSTNOT09), 1957981F6 (CONNNOT01), 3963055H1 (PROSNOT14), 5554170H1 (TONSDIT01), SASA02396F1, SASA00164F1

Table 2

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Genbank Homologs	Identification	Analytical Methods
1	693	S201 T373 S80 S227 S253 S391 T469 S674 S36 T41 S113 S197 S218 T368 T442 T603 S660 S678 S690 Y558	N39 N148 N171 N234 N303 N324 N341	Signal Peptide: M1-G25 Transmembrane Domain: F578-I596, Y641-M661, L610-F630 Targeting Signal: S691 TM7XN1 protein [Homo sapiens] (g4456467) Seven transmembrane receptor HE6 [Homo sapiens] (g2117161)	Seven-span transmembrane receptor	BLAST BLOCKS HMM MOTIFS PRINTS
2	84	S65 S36 T41 S51 S69 S81	N39	Signal Peptide: M1-G25 TM7XN1 protein [Homo sapiens] (g4456467) Seven transmembrane receptor HE6 [Homo sapiens] (g2117161)	Seven-span transmembrane receptor	BLAST HMM
3	136	T80 T124 T64 S86	N67	Asx Hydroxylation site: C41-C52 EGF-Like Domain: P22-K37, C35-Y56 CD97 [Homo sapiens] (g3342740)	EGF-TM7 receptors family	BLAST BLOCKS HMM MOTIFS PRINTS

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Genbank Homologs	Identification	Analytical Methods
4	307	S228 T48 S87 S93 T288 S297	N5	Transmembrane domain: I23-G41, I198-G41 Signal peptide: M1-S20 G-protein coupled receptors: G110-I126 Olfactory receptor: L59-R80, F177-D191, M237-G252, V271-L282, T288-F302 7-transmembrane receptor: G41-Y287 Olfactory receptor 4 [Gallus gallus] (g1246534)	Olfactory receptor	BLAST BLOCKS HMM MOTIFS PFAM PRINTS ProfileScan SPScan
5	318	S272 S57 S70 S111 S267	N8 N45 N91	Transmembrane domain: I34-V52, G202-L225 Signal peptide: M1-A26 G-protein coupled receptors: M113-I129 Olfactory receptor: M62-K83, S180-D194, F240-V255 7-transmembrane receptor: G44-Y294 Putative G-protein coupled receptor R1c [Rattus norvegicus] (g3420759)	Olfactory receptor	BLAST BLOCKS HMM MOTIFS PFAM PRINTS ProfileScan SPScan

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Genbank Homologs	Identification	Analytical Methods
6	138	T6 S22 S125	N44 N47 N83	mas product homolog modulating intracellular angiotensin II [Homo sapiens] (g244210)	MAS-like GPCR	BLAST
7	199	S21 S65 T93 S136 S155 T131		Transmembrane domain: L163-M184 Signal peptide: M1-G20 Rhodopsin-like GPCR superfamily: S160-M184	Rhodopsin-like GPCR	HMM PRINTS SPScan
8	445	S435 S273	N129	Transmembrane domain: V366-F386 Signal peptide: M1-A25 Contains similarity to G-coupled protein receptors [Caenorhabditis elegans] (g2736345)	GPCR	BLAST HMM SPScan

Table 3

Nucleotide SEQ ID NO:	Range of Useful Nucleotide Fragment(s)	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
9	1879-1928	Reproductive (0.290) Nervous (0.215) Gastrointestinal (0.131)	Cancer (0.495) Inflammation (0.168) Trauma (0.140)	pINCY
10		Nervous (0.318) Reproductive (0.227) Urologic (0.136)	Cancer (0.636) Inflammation (0.136) Trauma (0.091)	pINCY
11	325-387 667-732	Hematopoietic/Immune (0.625) Reproductive (0.250) Musculoskeletal (0.125)	Cancer (0.500) Inflammation (0.400)	pINCY
12	202-246	Nervous (1.000)	Cancer and cell proliferation (1.000)	PSPORT1
13	226-270	Reproductive (0.500) Urologic (0.500)	Cancer and cell proliferation (1.000)	PSPORT1
14	526-573	Gastrointestinal (0.500) Reproductive (0.500)	Cancer and cell proliferation (1.000)	PSPORT1
15	191-235	Developmental (0.167) Gastrointestinal (0.167) Hematopoietic/Immune (0.167) Musculoskeletal (0.167) Nervous (0.167) Reproductive (0.167)	Cancer and cell proliferation (0.834) Inflammation (0.167)	pINCY
16	802-846 1627-1672	Reproductive (0.290) Gastrointestinal (0.194) Nervous (0.161)	Cancer and cell proliferation (0.609) Inflammation (0.217)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Comment
9	BRAINOT14	Library was constructed using RNA isolated from brain tissue removed from the left frontal lobe of a 40-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology for the associated tumor tissue indicated grade 4 gemistocytic astrocytoma.
10	BRSTNOT12	Library was constructed using RNA isolated from diseased breast tissue removed from a 32-year-old Caucasian female during a bilateral reduction mammoplasty. Pathology indicated nonproliferative fibrocystic disease. Family history included benign hypertension and atherosclerotic coronary artery disease.
11	CONNTUT04	Library was constructed using RNA isolated from tumorous spinal tissue removed from a 35-year-old Caucasian male during an exploratory laparotomy. Pathology indicated schwannoma with degenerative changes. Patient history included anxiety, depression, neurofibromatosis and benign neoplasm of the scrotum. Previously the patient had a spinal fusion. Family history included brain cancer, liver disease, and multiple sclerosis.
12	BRAITUT03	Library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe a 17-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a grade 4 fibrillary giant and small-cell astrocytoma. Family history included benign hypertension and cerebrovascular disease.
13	KIDNTUT01	Library was constructed using RNA isolated from the kidney tumor tissue removed from an 8-month-old female during nephroureterectomy. Pathology indicated Wilms' tumor (nephroblastoma), which involved 90 percent of the renal parenchyma. Prior to surgery, the patient was receiving heparin anticoagulant therapy.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
14	PROSNOT06	Library was constructed using RNA isolated from the diseased prostate tissue of a 57-year-old Caucasian male during radical prostatectomy, removal of both testes and excision of regional lymph nodes. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated adenocarcinoma (Gleason grade 3+3). Patient history included a benign neoplasm of the large bowel and type I diabetes. Family history included a malignant neoplasm of the prostate and type I diabetes.
15	LIVRFET02	Library was constructed using RNA isolated from liver tissue removed from a Caucasian female fetus, who died at 20 weeks' gestation. Family history included seven days of erythromycin treatment for bronchitis in the mother during the first trimester.
16	TONSDIT01	Library was constructed using RNA isolated from the tonsil tissue of a 6-year-old Caucasian male during adenotonsillectomy. Pathology indicated lymphoid hyperplasia of the tonsils. The patient presented with an abscess of the pharynx. The patient was not taking any medications. Family history included hypothyroidism in the grandparent(s) and benign skin neoplasm in the sibling(s).

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) <i>J. Mol. Biol.</i> 215:403-410; Altschul, S.F. et al. (1997) <i>Nucleic Acids Res.</i> 25: 3389-3402.	EST's: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) <i>Proc. Natl. Acad. Sci.</i> 85:2444-2448; Pearson, W.R. (1990) <i>Methods Enzymol.</i> 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) <i>Adv. Appl. Math.</i> 2:482-489.	EST's: fasta E value=1.0E-6 Assembled EST's: fasta Identity= 95% or greater and Match length=200 bases or greater, fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS and PRINTS databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, <i>Nucl. Acid Res.</i> , 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) <i>Methods Enzymol.</i> 266:88-105; and Alwood, T.K. et al. (1997) <i>J. Chem. Inf. Comput. Sci.</i> 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and Probability value= 1.0E-3 or less if applicable
PFAM	A Hidden Markov Models-based application useful for protein family search.	Krogh, A. et al. (1994) <i>J. Mol. Biol.</i> , 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) <i>Nucleic Acids Res.</i> 26:320-322.	Score=10-50 bits, depending on individual protein families

Table 5 cont.

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Score= 4.0 or greater
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater, Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <u>supra</u> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and fragments thereof.
2. A substantially purified variant having at least 90% amino acid sequence identity to the amino acid sequence of claim 1.
3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.
4. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 3.
5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.
6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 3.
7. A method for detecting a polynucleotide, the method comprising the steps of:
 - (a) hybridizing the polynucleotide of claim 6 to at least one nucleic acid in a sample, thereby forming a hybridization complex; and
 - (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide in the sample.
8. The method of claim 7 further comprising amplifying the polynucleotide prior to hybridization.
9. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, and fragments thereof.
10. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 9.

11. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 9.
12. An expression vector comprising at least a fragment of the polynucleotide of claim 3.
- 5 13. A host cell comprising the expression vector of claim 12.
14. A method for producing a polypeptide, the method comprising the steps of:
- 10 a) culturing the host cell of claim 13 under conditions suitable for the expression of the polypeptide; and
- b) recovering the polypeptide from the host cell culture.
- 15 15. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.
16. A purified antibody which specifically binds to the polypeptide of claim 1.
17. A purified agonist of the polypeptide of claim 1.
- 20 18. A purified antagonist of the polypeptide of claim 1.
19. A method for treating or preventing a disorder associated with decreased expression or activity of GCRP, the method comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 15.
- 25 20. A method for treating or preventing a disorder associated with increased expression or activity of GCRP, the method comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 18.

1	M	-	-	-	-	-	-	-	-	T	P	Q	S	L	L	Q	T	T	L	F	L	L	S	L	-	-	-	L	F	L	V		2763296			
1	M	-	-	-	-	-	-	-	-	T	P	Q	S	L	L	Q	T	T	L	F	L	L	S	L	-	-	-	L	F	L	V		1595762			
1	M	V	F	S	V	R	Q	C	G	H	V	G	R	T	E	E	V	L	L	T	F	K	I	F	L	V	I	I	C	L	H	V	L	GI 2117161		
21	Q	G	A	H	G	R	G	H	R	E	D	F	R	-	-	-	-	-	-	-	F	C	S	Q	R	N	Q	T	H	R	S	S	2763296			
21	Q	G	A	H	G	R	G	H	R	E	D	F	R	-	-	-	-	-	-	-	F	C	S	Q	R	N	Q	T	H	R	S	S	1595762			
36	T	S	L	E	E	D	T	N	S	S	L	S	P	P	P	A	K	L	S	V	S	F	A	P	S	S	N	E	V	E	T	S	GI 2117161			
46	L	H	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2763296			
46	L	H	Y	K	P	T	P	D	L	R	I	S	I	E	N	S	E	E	A	L	T	V	H	A	P	P	A	A	H	P	A	S	R	-	1595762	
71	L	N	-	D	V	T	L	S	L	L	P	S	N	E	T	E	K	T	K	I	T	I	V	K	T	F	N	A	S	G	V	K	P	Q	GI 2117161	
49	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2763296			
80	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1595762			
105	N	I	C	N	L	S	S	I	C	N	D	S	A	F	F	R	G	E	I	M	F	Q	Y	D	K	E	S	T	V	P	Q	N	Q	H	I	GI 2117161
49	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2763296		
98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1595762		
140	T	N	G	T	L	T	G	V	L	S	L	S	E	L	K	R	S	E	L	N	K	T	L	Q	T	L	S	E	T	Y	F	I	M	C	A	GI 2117161
49	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2763296		
113	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1595762		
175	T	A	E	A	Q	S	T	L	N	C	T	F	T	I	K	L	N	N	T	M	N	A	C	A	A	I	A	A	L	E	R	V	K	I	R	GI 2117161

1/7

FIGURE 1A

FIGURE 1B

3/7

49 - - - - - 2763296
 233 I - - - - - 1595762
 420 L L K V V D D I G L Q L N F S N T T I S L T S P S L A L A V I R V N A G I 2117161

 49 - - - - - 2763296
 254 R Q - - - - - 1595762
 455 S S F N T T F V A Q D P A N L Q V S L E T Q A P E N S I G T I T L P G I 2117161

 49 - - - - - 2763296
 271 R T L F Q R T K G R S G E A E K R L L V D F S S Q A L F Q D K N S S 1595762
 490 S S L M N N L P A H D M E L A S R V Q F N F E T P A L F Q D P S L E G I 2117161

 49 - - - - - 2763296
 306 Q V - L G E K V L G I V V Q N T K V A N L T E P V V L T F Q H Q L Q P 1595762
 525 N L S L I S Y V I S S S V A N L T V R N L T R N V T V T L K H I N P S G I 2117161

 49 - - - - - 2763296
 340 K N - V T L Q C V F W V E D P T L S S P G H W S S A G C E T V - R R E 1595762
 560 Q D E L T V R C V F W D L G R N - G G R G G W S D N G C S V K D R R L G I 2117161

 49 - - - - - 2763296
 373 T Q T S C F C N H L T Y F A V L M - V S S V E V D A V H K H Y L S L L 1595762
 594 N E T I C T C S H L T S F G V L L D L S R T S V L P A Q M M A L T F I G I 2117161

FIGURE 1C

4/7

49	SYVGC	VVS	ALAC	LV	TI	AA	YLC	SR	VP	LP	CR	KP	RD	Y	2763296
407	TYIG	CG	LS	SIF	LS	VT	LV	TY	I	-	-	-	-	-	1595762
629	TYIG	CG	LS	SIF	LS	VT	LV	TY	I	-	-	-	-	-	GI 2117161
49	TIKV	HMN	LL	LA	VF	LL	DT	SF	LL	SE	PV	AL	TG	SE	AG
442	PSK	I	L	I	Q	L	CA	LL	LL	N	L	V	F	LL	DS
658	PSK	I	L	I	Q	L	CA	LL	LL	N	L	V	F	LL	DS
49	ASA	I	F	L	H	F	S	L	L	T	C	L	S	W	M
477	SV	A	V	F	L	H	F	S	L	L	T	C	L	S	W
693	SV	A	V	F	L	H	F	S	L	L	T	C	L	S	W
49	VP	G	Y	L	L	K	L	S	A	M	G	W	G	F	P
512	IR	K	Y	I	L	K	F	C	I	V	G	W	G	V	P
728	IR	K	Y	I	L	K	F	C	I	V	G	W	G	V	P
49	LA	V	H	R	T	P	E	G	V	I	Y	P	S	M	C
547	GS	Y	G	K	F	P	N	G	S	P	-	D	D	F	C
761	GS	Y	G	K	F	P	N	G	S	P	-	D	D	F	C
49	FL	F	N	M	A	M	L	A	T	M	V	V	Q	I	L
582	FL	L	N	V	S	M	F	I	V	V	L	V	Q	I	L
795	FL	L	N	V	S	M	F	I	V	V	L	V	Q	I	L

FIGURE 1D

6/7

1	M	G	V	S	A	L	T	E	L	T	-	-	-	-	S	L	G	G	-	-	-	-	D	S	L	S	C	-	-	-	-	3367641
1	M	G	G	R	V	F	L	A	F	C	V	W	L	T	L	P	G	A	E	T	Q	D	S	R	G	C	A	R	W	C	GI 3342740	
20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3367641	
31	P	Q	D	S	S	C	V	N	A	T	A	C	R	C	N	P	G	F	S	S	F	S	E	I	I	T	T	P	D	I	GI 3342740	
25	N	E	C	A	T	P	S	K	V	S	C	G	K	F	S	D	C	W	N	T	E	G	S	Y	D	C	V	C	S	P	3367641	
61	N	E	C	A	T	L	S	K	V	S	C	G	K	F	S	D	C	W	N	T	E	G	S	Y	D	C	V	C	S	P	GI 3342740	
55	G	Y	E	P	V	S	G	A	K	T	F	K	N	E	S	E	N	T	C	Q	G	K	N	H	P	T	S	S	D	F	3367641	
91	G	Y	E	P	V	S	G	A	K	T	F	K	N	E	S	E	N	T	C	Q	D	V	D	E	C	Q	Q	N	P	R	GI 3342740	
85	P	S	M	R	F	G	V	T	R	A	I	L	A	A	S	R	E	Q	G	L	G	-	-	-	-	-	-	-	-	-	3367641	
121	L	C	K	S	Y	G	T	C	V	N	T	L	G	S	Y	T	C	Q	C	L	P	G	F	K	D	V	N	E	C	T	GI 3342740	
106	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3367641		
151	S	G	Q	N	P	C	H	S	S	T	H	C	L	N	N	V	G	S	Y	Q	C	R	C	R	P	G	W	Q	P	I	GI 3342740	

FIGURE 2A

7/7

[illegible]

FIGURE 2B

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

TANG, Y. Tom
 YUE, Henry
 LAL, Preeti
 BANDMAN, Olga
 AU-YOUNG, Janice
 REDDY, Roopa
 CORLEY, Neil C.
 GUEGLER, Karl J.
 GORGONE, Gina A.
 BAUGHN, Mariah R.
 AZIMZAI, Yalda

<120> G-PROTEIN COUPLED RECEPTOR PROTEINS

<130> PF-0612 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/167,219; unassigned; 60/133,585

<151> 1998-10-06; 1998-10-06; 1999-05-11

<160> 18

<170> PERL Program

<210> 1

<211> 693

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1595762CD1

<400> 1

Met	Thr	Pro	Gln	Ser	Leu	Leu	Gln	Thr	Thr	Leu	Phe	Leu	Leu	Ser
1				5					10					15
Leu	Leu	Phe	Leu	Val	Gln	Gly	Ala	His	Gly	Arg	Gly	His	Arg	Glu
				20					25					30
Asp	Phe	Arg	Phe	Cys	Ser	Gln	Arg	Asn	Gln	Thr	His	Arg	Ser	Ser
				35					40					45
Leu	His	Tyr	Lys	Pro	Thr	Pro	Asp	Leu	Arg	Ile	Ser	Ile	Glu	Asn
				50					55					60
Ser	Glu	Glu	Ala	Leu	Thr	Val	His	Ala	Pro	Phe	Pro	Ala	Ala	His
				65					70					75
Pro	Ala	Ser	Arg	Ser	Phe	Pro	Asp	Pro	Arg	Gly	Leu	Tyr	His	Phe
				80					85					90
Cys	Leu	Tyr	Trp	Asn	Arg	His	Ala	Gly	Arg	Leu	His	Leu	Leu	Tyr
				95					100					105
Gly	Lys	Arg	Asp	Phe	Leu	Leu	Ser	Asp	Lys	Ala	Ser	Ser	Leu	Leu

	110		115	120
Cys Phe Gln His Gln Glu Glu Ser Leu	Ala Gln Gly Pro Pro Leu			
	125		130	135
Leu Ala Thr Ser Val Thr Ser Trp Trp	Ser Pro Gln Asn Ile Ser			
	140		145	150
Leu Pro Ser Ala Ala Ser Phe Thr Phe	Ser Phe His Ser Pro Pro			
	155		160	165
His Thr Ala Ala His Asn Ala Ser Val	Asp Met Cys Glu Leu Lys			
	170		175	180
Arg Asp Leu Gln Leu Leu Ser Gln Phe	Leu Lys His Pro Gln Lys			
	185		190	195
Ala Ser Arg Arg Pro Ser Ala Ala Pro	Ala Ser Gln Gln Leu Gln			
	200		205	210
Ser Leu Glu Ser Lys Leu Thr Ser Val	Arg Phe Met Gly Asp Met			
	215		220	225
Val Ser Phe Glu Glu Asp Arg Ile Asn	Ala Thr Val Trp Lys Leu			
	230		235	240
Gln Pro Thr Ala Gly Leu Gln Asp Leu	His Ile His Ser Arg Gln			
	245		250	255
Glu Glu Glu Gln Ser Glu Ile Met Glu	Tyr Ser Val Leu Leu Pro			
	260		265	270
Arg Thr Leu Phe Gln Arg Thr Lys Gly	Arg Ser Gly Glu Ala Glu			
	275		280	285
Lys Arg Leu Leu Leu Val Asp Phe Ser	Ser Gln Ala Leu Phe Gln			
	290		295	300
Asp Lys Asn Ser Ser Gln Val Leu Gly	Glu Lys Val Leu Gly Ile			
	305		310	315
Val Val Gln Asn Thr Lys Val Ala Asn	Leu Thr Glu Pro Val Val			
	320		325	330
Leu Thr Phe Gln His Gln Leu Gln Pro	Lys Asn Val Thr Leu Gln			
	335		340	345
Cys Val Phe Trp Val Glu Asp Pro Thr	Leu Ser Ser Pro Gly His			
	350		355	360
Trp Ser Ser Ala Gly Cys Glu Thr Val	Arg Arg Glu Thr Gln Thr			
	365		370	375
Ser Cys Phe Cys Asn His Leu Thr Tyr	Phe Ala Val Leu Met Val			
	380		385	390
Ser Ser Val Glu Val Asp Ala Val His	Lys His Tyr Leu Ser Leu			
	395		400	405
Leu Ser Tyr Val Gly Cys Val Val Ser	Ala Leu Ala Cys Leu Val			
	410		415	420
Thr Ile Ala Ala Tyr Leu Cys Ser Arg	Val Pro Leu Pro Cys Arg			
	425		430	435
Arg Lys Pro Arg Asp Tyr Thr Ile Lys	Val His Met Asn Leu Leu			
	440		445	450
Leu Ala Val Phe Leu Leu Asp Thr Ser	Phe Leu Leu Ser Glu Pro			
	455		460	465
Val Ala Leu Thr Gly Ser Glu Ala Gly	Cys Arg Ala Ser Ala Ile			
	470		475	480
Phe Leu His Phe Ser Leu Leu Thr Cys	Leu Ser Trp Met Gly Leu			
	485		490	495
Glu Gly Tyr Asn Leu Tyr Arg Leu Val	Val Glu Val Phe Gly Thr			
	500		505	510
Tyr Val Pro Gly Tyr Leu Leu Lys Leu	Ser Ala Met Gly Trp Gly			
	515		520	525

Phe	Pro	Ile	Phe	Leu	Val	Thr	Leu	Val	Ala	Leu	Val	Asp	Val	Asp
				530					535					540
Asn	Tyr	Gly	Pro	Ile	Ile	Leu	Ala	Val	His	Arg	Thr	Pro	Glu	Gly
				545					550					555
Val	Ile	Tyr	Pro	Ser	Met	Cys	Trp	Ile	Arg	Asp	Ser	Leu	Val	Ser
				560					565					570
Tyr	Ile	Thr	Asn	Leu	Gly	Leu	Phe	Ser	Leu	Val	Phe	Leu	Phe	Asn
				575					580					585
Met	Ala	Met	Leu	Ala	Thr	Met	Val	Val	Gln	Ile	Leu	Arg	Leu	Arg
				590					595					600
Pro	His	Thr	Gln	Lys	Trp	Ser	His	Val	Leu	Thr	Leu	Leu	Gly	Leu
				605					610					615
Ser	Leu	Val	Leu	Gly	Leu	Pro	Trp	Ala	Leu	Ile	Phe	Phe	Ser	Phe
				620					625					630
Ala	Ser	Gly	Thr	Phe	Gln	Leu	Val	Val	Leu	Tyr	Leu	Phe	Ser	Ile
				635					640					645
Ile	Thr	Ser	Phe	Gln	Gly	Phe	Leu	Ile	Phe	Ile	Trp	Tyr	Trp	Ser
				650					655					660
Met	Arg	Leu	Gln	Ala	Arg	Gly	Gly	Pro	Ser	Pro	Leu	Lys	Ser	Asn
				665					670					675
Ser	Asp	Ser	Ala	Arg	Leu	Pro	Ile	Ser	Ser	Gly	Ser	Thr	Ser	Ser
				680					685					690
Ser	Arg	Ile												

<210> 2

<211> 84

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2763296CD1

<400> 2

Met	Thr	Pro	Gln	Ser	Leu	Leu	Gln	Thr	Thr	Leu	Phe	Leu	Leu	Ser
1				5					10					15
Leu	Leu	Phe	Leu	Val	Gln	Gly	Ala	His	Gly	Arg	Gly	His	Arg	Glu
				20					25					30
Asp	Phe	Arg	Phe	Cys	Ser	Gln	Arg	Asn	Gln	Thr	His	Arg	Ser	Ser
				35					40					45
Leu	His	Tyr	Tyr	Trp	Ser	Met	Arg	Leu	Gln	Ala	Arg	Gly	Gly	Pro
				50					55					60
Ser	Pro	Leu	Lys	Ser	Asn	Ser	Asp	Ser	Ala	Arg	Leu	Pro	Ile	Ser
				65					70					75
Ser	Gly	Ser	Thr	Ser	Ser	Ser	Arg	Ile						
				80										

<210> 3

<211> 136

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3367641CD1

<400> 3

Met	Gly	Val	Ser	Ala	Leu	Thr	Glu	Leu	Thr	Ser	Leu	Gly	Gly	Asp
1				5					10					15
Ser	Leu	Ser	Cys	Cys	Val	Pro	Asp	Ile	Asn	Glu	Cys	Ala	Thr	Pro
			20						25					30
Ser	Lys	Val	Ser	Cys	Gly	Lys	Phe	Ser	Asp	Cys	Trp	Asn	Thr	Glu
			35						40					45
Gly	Ser	Tyr	Asp	Cys	Val	Cys	Ser	Pro	Gly	Tyr	Glu	Pro	Val	Ser
			50						55					60
Gly	Ala	Lys	Thr	Phe	Lys	Asn	Glu	Ser	Glu	Asn	Thr	Cys	Gln	Gly
			65						70					75
Lys	Asn	His	Pro	Thr	Ser	Ser	Asp	Phe	Pro	Ser	Met	Arg	Phe	Gly
			80						85					90
Val	Thr	Arg	Ala	Ile	Leu	Ala	Ala	Ser	Arg	Glu	Gln	Gly	Leu	Gly
			95						100					105
Cys	Arg	Val	Ser	Ala	Trp	Arg	Leu	Arg	Trp	Asp	Arg	Cys	Thr	Cys
			110						115					120
Thr	Tyr	Pro	Thr	Thr	Pro	Glu	Arg	Gln	Ser	Asp	Val	Arg	Ile	Lys
			125						130					135

Gly

<210> 4

<211> 307

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 866390CD1

<400> 4

Met	Glu	Ser	Glu	Asn	Arg	Thr	Val	Ile	Arg	Glu	Phe	Ile	Leu	Leu
1				5					10					15
Gly	Leu	Thr	Gln	Ser	Gln	Asp	Ile	Gln	Leu	Leu	Val	Phe	Val	Leu
			20						25					30
Val	Leu	Ile	Phe	Tyr	Phe	Ile	Ile	Leu	Pro	Gly	Asn	Phe	Leu	Ile
			35						40					45
Ile	Phe	Thr	Ile	Lys	Ser	Asp	Pro	Gly	Leu	Thr	Ala	Pro	Leu	Tyr
			50						55					60
Phe	Phe	Leu	Gly	Asn	Leu	Ala	Phe	Leu	Asp	Ala	Ser	Tyr	Ser	Phe
			65						70					75
Ile	Val	Ala	Pro	Arg	Met	Leu	Val	Asp	Phe	Leu	Ser	Ala	Lys	Lys
			80						85					90
Ile	Ile	Ser	Tyr	Arg	Gly	Cys	Ile	Thr	Gln	Leu	Phe	Phe	Leu	His
			95						100					105
Phe	Leu	Gly	Gly	Gly	Glu	Gly	Leu	Leu	Leu	Val	Val	Met	Ala	Phe
			110						115					120
Asp	Arg	Tyr	Ile	Ala	Ile	Cys	Arg	Pro	Leu	His	Tyr	Pro	Thr	Val
			125						130					135

Met Asn Pro Arg Thr Cys Tyr Ala Met Met Leu Ala Leu Trp Leu

	140		145		150
Gly Gly Phe Val His Ser Ile Ile Gln Val Val Leu Ile Leu Arg					
	155		160		165
Leu Pro Phe Cys Gly Pro Asn Gln Leu Asp Asn Phe Phe Cys Asp					
	170		175		180
Val Pro Gln Val Ile Lys Leu Ala Cys Thr Asp Thr Phe Val Val					
	185		190		195
Glu Leu Leu Met Val Phe Asn Ser Gly Leu Met Thr Leu Leu Cys					
	200		205		210
Phe Leu Gly Leu Leu Ala Ser Tyr Ala Val Ile Leu Cys Arg Ile					
	215		220		225
Arg Gly Ser Ser Ser Glu Ala Lys Asn Lys Ala Met Ser Thr Cys					
	230		235		240
Ile Thr His Ile Ile Val Ile Phe Phe Met Phe Gly Pro Gly Ile					
	245		250		255
Phe Ile Tyr Thr Arg Pro Phe Arg Ala Phe Pro Ala Asp Lys Val					
	260		265		270
Val Ser Leu Phe His Thr Val Ile Phe Pro Leu Leu Asn Pro Val					
	275		280		285
Ile Tyr Thr Leu Arg Asn Gln Glu Val Lys Ala Ser Met Lys Lys					
	290		295		300
Val Phe Asn Lys His Ile Ala					
	305				

<210> 5

<211> 318

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 998550CD1

<400> 5

Met Met Val Asp Pro Asn Gly Asn Glu Ser Ser Ala Thr Tyr Phe					
1	5		10		15
Ile Leu Ile Gly Leu Pro Gly Leu Glu Glu Ala Gln Phe Trp Leu					
	20		25		30
Ala Phe Pro Leu Cys Ser Leu Tyr Leu Ile Ala Val Leu Gly Asn					
	35		40		45
Leu Thr Ile Ile Tyr Ile Val Arg Thr Glu His Ser Leu His Glu					
	50		55		60
Pro Met Tyr Ile Phe Leu Cys Met Leu Ser Gly Ile Asp Ile Leu					
	65		70		75
Ile Ser Thr Ser Ser Met Pro Lys Met Leu Ala Ile Phe Trp Phe					
	80		85		90
Asn Ser Thr Thr Ile Gln Phe Asp Ala Cys Leu Leu Gln Met Phe					
	95		100		105
Ala Ile His Ser Leu Ser Gly Met Glu Ser Thr Val Leu Leu Ala					
	110		115		120
Met Ala Phe Asp Arg Tyr Val Ala Ile Cys His Pro Leu Arg His					
	125		130		135
Ala Thr Val Leu Thr Leu Pro Arg Val Thr Lys Ile Gly Val Ala					
	140		145		150

Ala Val Val Arg Gly Ala Ala Leu Met Ala Pro Leu Pro Val Phe		
155	160	165
Ile Lys Gln Leu Pro Phe Cys Arg Ser Asn Ile Leu Ser His Ser		
170	175	180
Tyr Cys Leu His Gln Asp Val Met Lys Leu Ala Cys Asp Asp Ile		
185	190	195
Arg Val Asn Val Val Tyr Gly Leu Ile Val Ile Ile Ser Ala Ile		
200	205	210
Gly Leu Asp Ser Leu Leu Ile Ser Phe Ser Tyr Leu Leu Ile Leu		
215	220	225
Lys Thr Val Leu Gly Leu Thr Arg Glu Ala Gln Ala Lys Ala Phe		
230	235	240
Gly Thr Cys Val Ser His Val Cys Ala Val Phe Ile Phe Tyr Val		
245	250	255
Pro Phe Ile Gly Leu Ser Met Val His Arg Phe Ser Lys Arg Arg		
260	265	270
Asp Ser Pro Leu Pro Val Ile Leu Ala Asn Ile Tyr Leu Leu Val		
275	280	285
Pro Pro Val Leu Asn Pro Ile Val Tyr Gly Val Lys Thr Lys Glu		
290	295	300
Ile Arg Gln Arg Ile Leu Arg Leu Phe His Val Ala Thr His Ala		
305	310	315
Ser Glu Pro		

<210> 6

<211> 138

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1547873CD1

<400> 6

Met Met Pro Thr Gln Thr Leu Pro Glu Thr His Ala Pro Gly Glu		
1	5	10
Thr Gln Ile Ser Leu Phe Ser Ser Gly Asp Gln Glu Ala Gln Asn		
20	25	30
Pro Asn Leu Val Ser Gln Leu Cys Gly Val Phe Leu Gln Asn Glu		
35	40	45
Thr Asn Glu Thr Ile His Met Gln Met Ser Met Ala Val Gly Gln		
50	55	60
Gln Ala Leu Pro Leu Asn Ile Ile Ala Pro Lys Ala Val Leu Val		
65	70	75
Ser Leu Cys Gly Val Leu Leu Asn Gly Thr Val Phe Trp Leu Leu		
80	85	90
Cys Cys Gly Ala Thr Asn Pro Tyr Met Val Tyr Ile Leu His Leu		
95	100	105
Val Ala Ala Asp Val Ile Tyr Leu Cys Cys Ser Ala Val Gly Phe		
110	115	120
Leu Gln Ile Tyr Ser Val Ala Glu Asp Met Val Ser Phe Val His		
125	130	135
His Gly Val		

<210> 7
 <211> 199
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2048484CD1

<400> 7
 Met Gly Leu Thr Leu Leu Leu Leu Leu Leu Gly Leu Glu Gly
 1 5 10 15
 Gln Gly Ile Val Gly Ser Leu Pro Glu Val Leu Gln Ala Pro Val
 20 25 30
 Gly Ser Ser Ile Leu Val Gln Cys His Tyr Arg Leu Gln Asp Val
 35 40 45
 Lys Ala Gln Lys Val Trp Cys Arg Phe Leu Pro Glu Gly Cys Gln
 50 55 60
 Pro Leu Val Ser Ser Ala Val Asp Arg Arg Ala Pro Ala Gly Arg
 65 70 75
 Arg Thr Phe Leu Thr Asp Leu Gly Gly Gly Leu Leu Gln Val Glu
 80 85 90
 Met Val Thr Leu Gln Glu Glu Asp Ala Gly Glu Tyr Gly Cys Met
 95 100 105

 Val Asp Gly Ala Arg Gly Pro Gln Ile Leu His Arg Val Ser Leu
 110 115 120
 Asn Ile Leu Pro Pro Glu Glu Glu Glu Glu Thr His Lys Ile Gly
 125 130 135
 Ser Leu Ala Glu Asn Ala Phe Ser Asp Pro Ala Gly Ser Ala Asn
 140 145 150
 Pro Leu Glu Pro Ser Gln Asp Glu Lys Ser Ile Pro Leu Ile Trp
 155 160 165
 Gly Ala Val Leu Leu Val Gly Leu Leu Val Ala Ala Val Val Leu
 170 175 180
 Phe Ala Val Met Ala Lys Arg Lys Gln Glu Ser Leu Leu Ser Gly
 185 190 195
 Pro Pro Arg Gln

<210> 8
 <211> 445
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5554170CD1

<400> 8
 Met Ala Ala Pro Thr Pro Ala Arg Pro Val Leu Thr His Leu Leu
 1 5 10 15
 Val Ala Leu Phe Gly Met Gly Ser Trp Ala Ala Val Asn Gly Ile
 20 25 30

Trp	Val	Glu	Leu	Pro	Val	Val	Val	Lys	Glu	Leu	Pro	Glu	Gly	Trp	35	40	45
Ser	Leu	Pro	Ser	Tyr	Val	Ser	Val	Leu	Val	Ala	Leu	Gly	Asn	Leu	50	55	60
Gly	Leu	Leu	Val	Val	Thr	Leu	Trp	Arg	Arg	Leu	Ala	Pro	Gly	Lys	65	70	75
Asp	Glu	Gln	Val	Pro	Ile	Arg	Val	Val	Gln	Val	Leu	Gly	Met	Val	80	85	90
Gly	Thr	Ala	Leu	Leu	Ala	Ser	Leu	Trp	His	His	Val	Ala	Pro	Val	95	100	105
Ala	Gly	Gln	Leu	His	Ser	Val	Ala	Phe	Leu	Ala	Leu	Ala	Phe	Val	110	115	120
Leu	Ala	Leu	Ala	Cys	Cys	Ala	Ser	Asn	Val	Thr	Phe	Leu	Pro	Phe	125	130	135
Leu	Ser	His	Leu	Pro	Pro	Arg	Phe	Leu	Arg	Ser	Phe	Phe	Leu	Gly	140	145	150
Gln	Gly	Leu	Ser	Ala	Leu	Leu	Pro	Cys	Val	Leu	Ala	Leu	Val	Gln	155	160	165
Gly	Val	Gly	Arg	Leu	Glu	Cys	Pro	Pro	Ala	Pro	Ile	Asn	Gly	Thr	170	175	180
Pro	Gly	Pro	Pro	Leu	Asp	Phe	Leu	Glu	Arg	Phe	Pro	Ala	Ser	Thr	185	190	195
Phe	Phe	Trp	Ala	Leu	Thr	Ala	Leu	Leu	Val	Ala	Ser	Ala	Ala	Ala	200	205	210
Phe	Gln	Gly	Leu	Leu	Leu	Leu	Pro	Pro	Pro	Pro	Pro	Ser	Val	Pro	215	220	225
Thr	Gly	Glu	Leu	Gly	Ser	Gly	Leu	Gln	Val	Gly	Ala	Pro	Gly	Ala	230	235	240
Glu	Glu	Glu	Val	Glu	Glu	Ser	Ser	Pro	Leu	Gln	Glu	Pro	Pro	Ser	245	250	255
Gln	Ala	Ala	Gly	Thr	Thr	Pro	Gly	Pro	Asp	Pro	Lys	Ala	Tyr	Gln	260	265	270
Leu	Leu	Ser	Ala	Arg	Ser	Ala	Cys	Leu	Leu	Gly	Leu	Leu	Ala	Ala	275	280	285
Thr	Asn	Ala	Leu	Thr	Asn	Gly	Val	Leu	Pro	Ala	Val	Gln	Ser	Phe	290	295	300
Ser	Cys	Leu	Pro	Tyr	Gly	Arg	Leu	Ala	Tyr	His	Leu	Ala	Val	Val	305	310	315
Leu	Gly	Ser	Ala	Ala	Asn	Pro	Leu	Ala	Cys	Phe	Leu	Ala	Met	Gly	320	325	330
Val	Leu	Cys	Arg	Ser	Leu	Ala	Gly	Leu	Gly	Gly	Leu	Ser	Leu	Leu	335	340	345
Gly	Val	Phe	Cys	Gly	Gly	Tyr	Leu	Met	Ala	Leu	Ala	Val	Leu	Ser	350	355	360
Pro	Cys	Pro	Pro	Leu	Val	Gly	Thr	Ser	Ala	Gly	Val	Val	Leu	Val	365	370	375
Val	Leu	Ser	Trp	Val	Leu	Cys	Leu	Gly	Val	Phe	Ser	Tyr	Val	Lys	380	385	390
Val	Ala	Ala	Ser	Ser	Leu	Leu	His	Gly	Gly	Gly	Arg	Pro	Ala	Leu	395	400	405
Leu	Ala	Ala	Gly	Val	Ala	Ile	Gln	Val	Gly	Ser	Leu	Leu	Gly	Ala	410	415	420
Val	Ala	Met	Phe	Pro	Pro	Thr	Ser	Ile	Tyr	His	Val	Phe	His	Ser	425	430	435

Arg Lys Asp Cys Ala Asp Pro Cys Asp Ser
440 445

<210> 9
<211> 3632
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1595762CB1

<400> 9
ggaaggggag gagcaggcca cacaggcaca ggccggtgag ggacctgccc agacctggag 60
gtggtgactt ccaagagtga ctccgtcggg ggaatatgac tccccagtcg ctgctgcaga 120
cgacactgtt cctgctgagt ctgctcttcc tggccaagg tgcccacggc agggggccaca 180
gggaagactt tcgcttctgc agccagcggg accagacaca caggagcagc ctccactaca 240
aaccacacac agacctgcgc atctccatcg agaactccga agaggccctc acagtccatg 300
cccccttccc tgcagcccac cctgcttccc gatccttccc tgaccccagg ggctcttacc 360
acttctgcct ctactggaac cgacatgctg ggagattaca tcttctctat ggcaagcgtg 420
acttcttgct gagtgcacaa gcctctagcc tctctgctt ccagcaccag gaggagagcc 480
tggctcaggg ccccccgtg ttagccactt ctgtcacctc ctggtggagc cctcagaaca 540
tcagcctgcc cagtgcggcc agcttcacct tctccttcca cagtccctcc cacaaggccg 600
ctcacaatgc ctcggtggac atgtgcgagc tcaaaaggga cctccagctg ctccagccagt 660
tctgaagca tccccagaag gcctcaagga ggccctcggc tgcccccgcc agccagcagt 720
tgcagagcct ggagtcgaaa ctgacctctg tgagattcat gggggacatg gtgtccttcg 780
aggaggaccg gatcaacgcc acggtgtgga agctccagcc cacagccggc ctccaggacc 840
tgcacatcca ctcccgccag gaggaggagc agagcgagat catggagtac tcggtgctgc 900
tgcctcgaac actcttccag aggacgaaag gccggagcgg ggaggctgag aagagactcc 960
tctggtgga ctccagcagc caagccctgt tccaggacaa gaattccagc caagtcctgg 1020
gtgagaaggc cttggggatt gtggtacaga acaccaaagt agccaacctc acggagcccc 1080
tgggtgtcac ctccagcac cagctacagc cgaagaatgt gactctgcaa tgtgtgttct 1140
gggttgaaga cccacattg agcagcccg ggcatggag cagtgtggg tgtgagaccg 1200
tcaggagaga aaccacaaac tctgtcttct gcaaccactt gacctacttt gcagtgtcta 1260
tgggtctcctc ggtggaagtg gacgccgtgc acaagcacta cctgagcctc ctctcctacg 1320
tgggctgtgt cgtctctgcc ctggcctgcc ttgtcaccat tgccgcctac ctctgctcca 1380
gggtgcccc cccgtgcagg aggaacctc gggactacac catcaagggtg cacatgaacc 1440
tgctgctggc cgtcttctct ctggacacga gcttctgct cagcgagccg gtggccctga 1500
caggctctga ggctggctgc cgagccagt ccacttctct gcacttctcc ctgtcacct 1560
gccttctctg gatgggcctc gaggggtaca acctctaccg actcgtgggtg gaggtctttg 1620
gcacctatgt ccctggctac ctactcaagc tgagcgccat gggctggggc ttccccatct 1680
ttctggtgac gctggtggcc ctggtggatg tggacaacta tggcccatc atcttggtg 1740
tgcataggac tccagagggc gtcacttacc ctccatgtg ctggatccgg gactccctgg 1800
tcagctacat caccaacctg ggctcttca gcctggtgtt tctgttcaac atggccatgc 1860
tagccaccat ggtggtgcag atcctgcggc tgcgccccca caccacaaag tggtcacatg 1920
tgctgacact gctgggcctc agcctggtcc ttggcctgcc ctgggccttg atcttcttct 1980
ccttggcttc tggcaccttc cagcttctgc tctctacct ttccagcatc atcacctcct 2040
tccaaggctt cctcatcttc atctggtact ggtccatgcg gctgcaggcc cggggtggcc 2100
cctccccctt gaagagcaac tcagacagcg ccaggctccc catcagctcg ggcagcacct 2160
cgtccagccg catctaggcc tccagcccac ctgcccattg gatgaagcag agatgcggcc 2220
tcgtcgaca ctgctgtgg ccccgagcc cggccagcc ccaggccagt cagccgcaga 2280
ctttggaaag cccaacgacc atggagagat gggccgttgc catggtggac ggactcccg 2340
gctgggcttt tgaattggcc ttggggacta ctgggtctc actcagctcc caggggactc 2400
agaagtgcgc cgccatgctg cctagggtac tgtccccaca tctgtcccaa ccagctgga 2460

```

ggcctgggtct ctccttataa cccctggggc cagccctcat tgctgggggc caggccttgg 2520
atcttgaggg tctggcacat ccttaatect gtgcccctgc ctgggacaga aatgtggctc 2580
cagttgtctct gtctctctgt gtcacctga gggcactctg catctctgt cattttaacc 2640
tcagggtggca cccagggcga atggggccca gggcagacct tcagggccag agccctggcg 2700
gaggagagggc cctttgccag gagcacagca gcagctcgcc tacctctgag cccaggcccc 2760
ctccctccct cagcccccca gtccctccctc catcttccct ggggttctcc tcctctccca 2820
gggcctcctt gctccttctg tcacagctgg gggccccga ttccaatgct gttttttggg 2880
gagtggtttc caggagctgc ctggtgtctg ctgtaaagt ttgtctactg cacaagctc 2940
ggcctgcccc tgagccaggc tcggtaccga tgcgtgggt gggctaggct cctctgtcca 3000
tctgggcctt tgtatgagct gcattgccct tgctcaccct gaccaagcac acgcctcaga 3060
ggggccctca gcctctctg aagccctctt gtggcaagaa ctgtggacca tgccagtccc 3120
gtctgggttc catcccacca ctccaaggac tgagactgac ctctctggt gacactggcc 3180
tagagcctga cactctccta agaggttctc tccaagcccc caaatagctc caggcgccct 3240
cgcccgccca tcatggttaa ttctgtccaa caaacacaca cggttagatt gctggcctgt 3300
tgtagggtgg agggacacag atgaccgacc tggctactcc tcctgccaac attcagtctg 3360
gtatgtgagg cgtgcgtgaa gcaagaactc ctggagctac agggacaggg agccatcatt 3420
cctgcctggg aatcctggaa gacttctctg aggagtcagc gttcaatctt gaccttgaag 3480
atgggaagga tggtcttttt acgtaccaat tcttttgtct tttgatatta aaaagaagta 3540
catgttcatt gtagagaatt tggaaactgt agaagagaat caagaagaaa aataaaaaatc 3600
agctgttgta atcacctagc aaaaaaaaaa aa 3632

```

<210> 10

<211> 714

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2763296CB1

<400> 10

```

gggagcctcc cagctctctc agctcactcg gcaggcagcg gggaccaggg ctggcaggtt 60
aagcctctgg ggggtgatcc tgaaagggtg tccagccgcc tggccctgcg tgggaccctc 120
cacctggcag caggtgggtga cttccaagag tgactccgtc ggaggaaaat gactccccag 180
tcgctgctgc agacgacact gttcctgctg agtctgctct tcctgggccca aggtgcccac 240
ggcagggggc acagggaaga ctttcgcttc tgcagccagc ggaaccagac acacaggagc 300
agcctccact actactggtc catgcggctg caggcccggt gtggccctc ccctctgaag 360
agcaactcag acagcgccag gctccccatc agctcgggca gcacctcgtc cagccgcctc 420
taggcctcca gccacactgc ccatgtaatg aagcagagat gcggcctcgt cgcacactgc 480
ctgtagcccc cgaaccggtc ccagccccag gccagtaagc cgcagacttt agaaagccca 540
acgaccatgg agagatgggc cgttgccatg gtggacggac tcccgggctg ggcttttgag 600
attggcttag gggctactcg gctctcactc agctcccacg ggactcaaga atgcggcgcc 660
atgctgcctt aggtactgtc cccacatctg tcccaaccca gctggaggcc tgggt 714

```

<210> 11

<211> 1035

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 1029

<223> a or g or c or t, unknown, or other

<220>
<221> misc_feature
<223> Incyte ID No: 3367641CB1

<400> 11
agactgggtct tgagctcctg agctcaggtg atccacctgc cttagcctcc caaaatgctg 60
ggattacagg cgtgagccac tgcacccggc cgccatgggg tttaaatttc taaaaagaga 120
ctgggggatgg ggatgtggag cttcctgtcc atccctcacc tcagcactgt tctgtgtgc 180
caaactctaag atgccagcca gagaactaag atcaagtgtg gtcttcagga atggcttgct 240
ccagctgcag catagcctgt gcagtgcac ttaagatttt ttccaaaaga tactaagcag 300
caggcccat gttggggagt caggatgggg gtctctgccc tgacagaact cacatctttg 360
ggaggtgact cctgtcctg ttgtgttcca gacatcaacg agtgtgcaac accgtcgaaa 420
gtgtcatgcy gaaaattctc ggactgctgg aacacagagg ggagctacga ctgctgtgc 480
agcccgggat atgagcctgt ttctggggca aaaacattca agaagtagag cgagaacacc 540
tgtcaaggta agaaccaccc cagctcctct gactttccat ccatgaggtt tggggtcacc 600
agagccattc tggcagcatc cagagagcag ggccttggtt gtagggtcag tgcctggcgt 660
ctgagatggg acaggtgcac atgtacctac cccaccaccc cagagaggca gagtgtgtg 720
aggattaagg gctgactctg ggaccagctg cctgggcca atcctgataa ccaagtgtat 780
gacgttggcc aggatactta ttcagaaacc gctgtccaca tgtgagatgt gataacgtta 840
tcatgcttgt aattccagtg ctttgggagg ctgaggcggg aggatctctt gaggccagga 900
gttcaagacc agcccatagc aacatagcaa gacccaccc ctataaaagg ggaggcgaag 960
gcaggtagat tgtttgagcc caggggttcg agaccagcat gagcaagcgg gtgaaacca 1020
tattattgng ttgca 1035

<210> 12
<211> 1002
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 866390CB1

<400> 12
atctggtaga actcagctgt gaattcctgt gatcctgggc tttttctggc cagggaaatg 60
gaaagcgaga acagaacagt gataagagaa ttcactctcc ttggtctgac ccagtctcaa 120
gatattcagc tctgtgtctt tgtgctagt ttaatatct acttcatcat cctccctgga 180
aattttctca ttattttcac cataaagtca gacctgggc tcacagcccc cctctatttc 240
tttctgggca acttggcctt cctggatgca tctactcct tcattgtggc tccccggatg 300
ttggtggact tctctctgc gaagaagata atctcctaca gaggtgcat cactcagctc 360
ttttctctgc acttccctgg aggaggggag ggattactcc ttgttgtgat ggcctttgac 420
cgctacatcy ccatctgccg gcctctgcac tatcctactg tcatgaaccc tagaacctgc 480
tatgcaatga tgttggtctt gtggcttggg ggttttgtcc actccattat ccaggtggtc 540
ctcatcctcc gcttgccctt ttgtggccca aaccagctgg acaacttctt ctgtgatgtc 600
ccacaggtca tcaagctggc ctgcaccgac acatttgtgg tggagcttct gatggtcttc 660
aacagtggcc tgatgacact cctgtgcttt ctggggcttc tggcctccta tgcagtcatt 720
ctttgtcgca tacgagggtc ttcttctgag gcaaaaaaca aggcctatgtc cagctgcac 780
accatataca ttgttatatt cttcatgttt ggacctggca tcttcatcta cagcgcccc 840
ttcagggtt tccagctga caaggtggtt tctctcttcc acacagtgt ttttctttg 900
ttgaatcctg tcatattatac ccttcgcaac caggaagtga aagcttccat gaaaaagggtg 960
tttaataagc acatagcctg aaaaaggctg caaaaaaaaa aa 1002

<210> 13
<211> 1302
<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 998550CB1

<400> 13

```

tggacaaagg gggtcacaca ttcccttccat acggttgagc ctctacctgc ctggtgctgg 60
tcacagttca gcttcttcat gatggtggat cccaatggca atgaatccag tgctacatac 120
ttcatcctaa taggcctccc tggtttagaa gaggtcagc tctggttggc cttcccattg 180
tgctccctct acctattgc tgtgctaggt aacttgacaa tcatctacat tgtgcggaac 240
gagcacagcc tgcattgagc catgtatata ttcttttga tgctttcagg cattgacatc 300
ctcatctcca cctcatccat gcccaaatg ctggccatct tctggttcaa ttccactacc 360
atccagtttg atgcttgtct gctacagatg tttgccatcc actccttacc tggcatggaa 420
tccacagtgc tgcaggccat ggcttttgac cgtatgtgg ccatctgtca cccactgcgc 480
catgccacag tacttacgtt gcctcgtgct accaaaattg gtgtggctgc tgtggtgcgg 540
ggggctgcac tgatggcacc ccttcctgct tcatcaagc agctgccctt ctgccgctcc 600
aatatccttt cccattccta ctgcctacac caagatgtca tgaagctggc ctgtgatgat 660
atccgggtca atgtcgtcta tggccttacc gtcacatct ccgccattgg cctggactca 720
cttctcatct ccttctcata tctgcttatt cttaagactg tgttgggctt gacacgtgaa 780
gcccaggcca aggcatttgg cacttgctgc tctcatgtgt gtgctgtgtt catattctat 840
gtacctttta ttgattgtc catggtgcat cgcttttagc agcggcgtga ctctcgtg 900
ccgctcatct tggccaatat ctatctgctg gttcctcctg tgctcaaccc aattgtctat 960
ggagtgaaga caaaggagat tgcacagcgc atccttcgac ttttccatgt ggccacacac 1020
gcttcagagc cctaggtgtc agtgatcaaa cttcttttcc attcagagtc ctctgattca 1080
gattttaatg ttaacatttt ggaagacagt attcagaaaa aaaatttcct taataaaaaat 1140
acaactcaga tcttcaaat atgaaactgg ttggggaatc tccatttttt caatattatt 1200
ttcttctttg ttttcttgc acatataatt attaataccc tgactagggt gtggtttgag 1260
ggttattact tttcatttta ccatgcagtc caaatctaaa ct 1302

```

<210> 14

<211> 1136

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1547873CB1

<400> 14

```

aagagagagc tcatattcag aggacaggaa gacaattcca gaccattggt ccacagaggt 60
gtcacctgtc cttctgtgtc tgtgtcagcc tcaactgccc ctgacttacc cctgccccca 120
ggaaaaatgt cacagctgac atttgagca tagcagtcga tgatgccac acagacactg 180
cctgagactc acgcccctgg agaaacgcag atttccctat ttccagtggt tgaccaggag 240
gcacagaacc caaacctggt atctcagctc tgtggcgtct ttcttcaaaa tgagacgaat 300
gaaaccatac atatgcagat gagcatggca gtgggacagc aggcctgccc cttgaatata 360
attgccccca aggtgtgtct ggtctccctc tgtggggtct tattgaatgg cactgtcttc 420
tggctgcttt gctgtggggc cacgaatccc tacatggtat acatcctcca cctggctgct 480
gctgacgtga tctatctttg ctgctcggca gtggggttct tacagattta ctctgtggct 540
gaagacatgg tcagttttgt ccaccatggt gtttaaaata atgtatagtc tctttttaag 600
gtgtaatttg tcaataaact ttttttaatg atttctatat gtgcttattc aattatgtgt 660
acatgttact caaaacaaag aagattcatt caagtctgat attaaaatat atggttacct 720
cttaattccc atactgcac attgtgtagc tcatgtctca tgatgctgac atttcaggag 780
catgttaagg aggtgtcaga gtgaaggaa gggcatgaga cttctcgtgg acaggctctt 840
gtctctccac ttctgttagc tctccatcta taagcagatt agtccttttt cagttgtctg 900

```

```

agctgcagat acaatttcca acattttcat tggtcactta ctteggttat gattttaaaa 960
acatgttatt gtcaagcaca ccgtttttat aaatttgaat ttgtcagtct tttatggctt 1020
atgtattgag ttatgattaa agaagtctta cctacatgag ataataataa atgaatctat 1080
gctgtcttat actattaaat tatttcattt gtaaacaatgt aaaagaaaaa aaaaaa 1136

```

<210> 15

<211> 1185

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2048484CB1

<400> 15

```

tgcacagtg cccaggcaag cccaggagtt gacatttctc tgcccagcca tgggcctcac 60
cctgctcttg ctgctgctcc tgggactaga aggtcagggc atagttggca gcctccctga 120
ggtgctgcag gcacccgtgg gaagctccat tctggtgcag tgccactaca ggctccagga 180
tgtcaaagct cagaaggtgt ggtgccggtt cttgccggag ggggtgccagc ccctgggtgc 240
ctcagctgtg gatcgagag ctccagcggg caggcgtagc tttctcacag acctgggtgg 300
gggcctgctg caggtggaaa tggttaccct gcaggaagag gatgctggcg agtatggctg 360
catggtggat ggggccaggg ggcccagat tttgcacaga gtctctctga acatactgcc 420
cccagaggaa gaagaagaga ccataagat tggcagctctg gctgagaacg cattctcaga 480
ccctgcaggc agtgccaaacc ctttgggaacc cagccaggat gagaagagca tccccttgat 540
ctggggtgct gtgctcctgg taggtctgct ggtggcagcg gtggtgctgt ttgctgtgat 600
ggccaagagg aaacaagaat ccctcctcag tggtcacca cgtcagtgac tctggaccgg 660
ctgctgaatt gcctttggat gtaccacaca ttaggcttga ctcaccacct tcatttgaca 720
ataccaccta caccagccta cctcttgatt ccccatcagg aaaaccttca ctcccagctc 780
catcctcatt gccccctcta cctcctaagg tcctggtctg ctccaagcct gtgacatatg 840
ccacagtaat cttcccggga gggaacaagg gtggagggac ctctgttggg ccagcccaga 900
atccacctaa caatcagact ccatccagct aagctgctca tcacacttta aactcatgag 960
gaccatccct aggggttctg tgcacccatc cagccagctc atgccctagg atccttagga 1020
tatctgagca accagggact ttaagatcta atccaatgtc ctaactttac tagggaaagt 1080
gacgctcaga catgactgag atgtcttggg gaagacctcc ctgcacccaa ctccccact 1140
ggttcttcta ccattacaca ctgggctaaa taaaccctaa taatg 1185

```

<210> 16

<211> 2171

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5554170CB1

<400> 16

```

cccacgcgtc cgcccacgcg tccgcggggt cctgcgcgct ccggactgag gtggcgctcc 60
tgggccggac ggcggtgtcc cggcggtggcg ggaagccggc actggagcgg gagcgactg 120
ggcgcgggac cgggagggcg agggaccgga cggctcccga gtcgcccacc tgacggtacc 180
gagagggcgg cgcccctccg agcagagccg tcccggccac tcccctggga tctgacttgg 240
ctcttgcggt cgcgggcacc gtgaagccct ggggtgtgctg tggctcctcc tggtaggcgc 300
cctttcccgg cgtccggctt ggggtgtgtg tggcggtgac tccagccccg cctctccctg 360
gagaggaggg ctccactcgc tccttcggcc tcctccccctg gggccgcagc gactcggggc 420
ggcttctctg ttcctgcct gccggcggtc ccgctggcta gaagaagtct tcacttccca 480
ggagagccaa agcgtgtctg gccctaggtg ggaaaagaac tggctgtgac ctttgcctctg 540

```

```

acctggaagg gccagcctt gggtgaatg gcagaccca cgcgcgcgcg tccggtgctg 600
accacactgc tgggtgctct cttcggtcatg ggctcctggg ctgcggtcaa tgggatctgg 660
gtggagctac ctgtggtggt caaagagctt ccagagggtt ggagcctccc ctcttacgtc 720
tctgtgcttg tggctctggg gaacctgggt ctgctggtgg tgacctctg gaggaggctg 780
gccccaggaa aggacgagca ggtcccatc cgggtggtgc aggtgctggg catggtgggc 840
acagccctgc tggcctctct gtggcaccat gtggccccag tggcaggaca gttgcattct 900
gtggccttct tagcactggc ctttgtgctg gcactggcat gctgtgcctc gaatgtcact 960
ttcctgccct tcttgagcca cctgccacct cgcttcttac ggtcattctt cctgggtcaa 1020
ggcctgagtg ccctgctgcc ctgctgctg gccctagtgc aggtgtggtg ccgcctcgag 1080
tgcccgccag ccccatcaa cggcaccctt ggccccccgc tcgacttcct tgagcgtttt 1140
ccgcccagca ccttcttctg ggcactgact gcccttctgg tcgcttcagc tgctgccttc 1200
cagggctctt tgtgtgctgt gccgccacca ccatctgtac ccacagggga gttaggatca 1260
ggcctccagg tgggagcccc aggagcagag gaagagggtg aagagtcctc accactgcaa 1320
gagccaccaa gccaggcagc aggcaccacc cctgggtccag accctaaggc ctatcagctt 1380
ctatcagccc gcagtgcctg cctgctgggc ctggtggccg ccaccaacgc gctgaccaat 1440
ggcgtgctgc ctgcctgca gagcttttcc tgcttaccct acgggcgtct ggcctaccac 1500
ctggctgtgg tgctgggcag tgctgccaat cccctggcct gcttctctgg catgggtgtg 1560
ctgtgcaggt ccttggcagg gctgggcggc ctctctctgc tgggcgtgtt ctgtgggggc 1620
tacctgatgg cgctggcagt cctgagcccc tgcccgcccc tgggtgggcac ctgcgcgggg 1680
gtggtcctcg tgggtgctgc gtgggtgctg tgtcttggtg tgttctccta cgtgaagggtg 1740
gcagccagct ccctgctgca tggcgggggc cggccggcat tgctggcagc cggcgtggcc 1800
atccagggtg gctctctgct cggcgctgtt gctatgttcc ccccgaccag catctatcac 1860
gtgttcaca gcagaaagga ctgtgcagac cctgtgact cctgagcctg ggcagggtgg 1920
gaccccgctc cccaacacct gtctttccct caatgctgcc accatgcctg agtgcctgca 1980
gcccaggagg cccgcacacc ggtacactcg tggacaccta cacactccat aggagatcct 2040
ggctttccag ggtgggcaag ggcaaggagc aggcttggag ccagggacca gtgggggctg 2100
tagggtaagc ccctgagcct gggacctaca tgtggtttgc gtaataaaac atttgtattt 2160
aaaaaaaaa a 2171

```

<210> 17

<211> 1014

<212> PRT

<213> Homo sapiens

<300>

<308> GenBank ID No: g2117161

<400> 17

```

Met Val Phe Ser Val Arg Gln Cys Gly His Val Gly Arg Thr Glu
  1             5             10             15
Glu Val Leu Leu Thr Phe Lys Ile Phe Leu Val Ile Ile Cys Leu
      20             25             30
His Val Val Leu Val Thr Ser Leu Glu Glu Asp Thr Asp Asn Ser
      35             40             45
Ser Leu Ser Pro Pro Ala Lys Leu Ser Val Val Ser Phe Ala
      50             55             60
Pro Ser Ser Asn Glu Val Glu Thr Thr Ser Leu Asn Asp Val Thr
      65             70             75
Leu Ser Leu Leu Pro Ser Asn Glu Thr Glu Lys Thr Lys Ile Thr
      80             85             90
Ile Val Lys Thr Phe Asn Ala Ser Gly Val Lys Pro Gln Arg Asn
      95             100            105
Ile Cys Asn Leu Ser Ser Ile Cys Asn Asp Ser Ala Phe Phe Arg
      110            115            120

```


15/18

530	535	540
Thr Val Arg Asn Leu Thr Arg Asn Val	Thr Val Thr Leu Lys His	
545	550	555
Ile Asn Pro Ser Gln Asp Glu Leu Thr	Val Arg Cys Val Phe Trp	
560	565	570
Asp Leu Gly Arg Asn Gly Gly Arg Gly	Gly Trp Ser Asp Asn Gly	
575	580	585
Cys Ser Val Lys Asp Arg Arg Leu Asn	Glu Thr Ile Cys Thr Cys	
590	595	600
Ser His Leu Thr Ser Phe Gly Val Leu	Leu Asp Leu Ser Arg Thr	
605	610	615
Ser Val Leu Pro Ala Gln Met Met Ala	Leu Thr Phe Ile Thr Tyr	
620	625	630
Ile Gly Cys Gly Leu Ser Ser Ile Phe	Leu Ser Val Thr Leu Val	
635	640	645
Thr Tyr Ile Ala Phe Glu Lys Ile Arg	Arg Asp Tyr Pro Ser Lys	
650	655	660
Ile Leu Ile Gln Leu Cys Ala Ala Leu	Leu Leu Leu Asn Leu Val	
665	670	675
Phe Leu Leu Asp Ser Trp Ile Ala Leu	Tyr Lys Met Gln Gly Leu	
680	685	690
Cys Ile Ser Val Ala Val Phe Leu His	Tyr Phe Leu Leu Val Ser	
695	700	705
Phe Thr Trp Met Gly Leu Glu Ala Phe	His Met Tyr Leu Ala Leu	
710	715	720
Val Lys Val Phe Asn Thr Tyr Ile Arg	Lys Tyr Ile Leu Lys Phe	
725	730	735
Cys Ile Val Gly Trp Gly Val Pro Ala	Val Val Val Thr Ile Ile	
740	745	750
Leu Thr Ile Ser Pro Asp Asn Tyr Gly	Leu Gly Ser Tyr Gly Lys	
755	760	765
Phe Pro Asn Gly Ser Pro Asp Asp Phe	Cys Trp Ile Asn Asn Asn	
770	775	780
Ala Val Phe Tyr Ile Thr Val Val Gly	Tyr Phe Cys Val Ile Phe	
785	790	795
Leu Leu Asn Val Ser Met Phe Ile Val	Val Leu Val Gln Leu Cys	
800	805	810
Arg Ile Lys Lys Lys Lys Gln Leu Gly	Ala Gln Arg Lys Thr Ser	
815	820	825
Ile Gln Asp Leu Arg Ser Ile Ala Gly	Leu Thr Phe Leu Leu Gly	
830	835	840
Ile Thr Trp Gly Phe Ala Phe Phe Ala	Trp Gly Pro Val Asn Val	
845	850	855
Thr Phe Met Tyr Leu Phe Ala Ile Phe	Asn Thr Leu Gln Gly Phe	
860	865	870
Phe Ile Phe Ile Phe Tyr Cys Val Ala	Lys Glu Asn Val Arg Lys	
875	880	885
Gln Trp Arg Arg Tyr Leu Cys Cys Gly	Lys Leu Arg Leu Ala Glu	
890	895	900
Asn Ser Asp Trp Ser Lys Thr Ala Thr	Asn Gly Leu Lys Lys Gln	
905	910	915
Thr Val Asn Gln Gly Val Ser Ser Ser	Ser Asn Ser Leu Gln Ser	
920	925	930
Ser Ser Asn Ser Thr Asn Ser Thr Thr	Leu Leu Val Asn Asn Asp	
935	940	945

Cys Ser Val His Ala Ser Gly Asn Gly Asn Ala Ser Thr Glu Arg
 950 955 960
 Asn Gly Val Ser Phe Ser Val Gln Asn Gly Asp Val Cys Leu His
 965 970 975
 Asp Phe Thr Gly Lys Gln His Met Phe Asn Glu Lys Glu Asp Ser
 980 985 990
 Cys Asn Gly Lys Gly Arg Met Ala Leu Arg Arg Thr Ser Lys Arg
 995 1000 1005
 Gly Ser Leu His Phe Ile Glu Gln Met
 1010

<210> 18
 <211> 287
 <212> PRT
 <213> Homo sapiens

<300>
 <308> GenBank ID No: g3342740

<400> 18
 Met Gly Gly Arg Val Phe Leu Ala Phe Cys Val Trp Leu Thr Leu
 1 5 10 15
 Pro Gly Ala Glu Thr Gln Asp Ser Arg Gly Cys Ala Arg Trp Cys
 20 25 30
 Pro Gln Asp Ser Ser Cys Val Asn Ala Thr Ala Cys Arg Cys Asn
 35 40 45
 Pro Gly Phe Ser Ser Phe Ser Glu Ile Ile Thr Thr Pro Asp Ile
 50 55 60
 Asn Glu Cys Ala Thr Leu Ser Lys Val Ser Cys Gly Lys Phe Ser
 65 70 75
 Asp Cys Trp Asn Thr Glu Gly Ser Tyr Asp Cys Val Cys Ser Pro
 80 85 90
 Gly Tyr Glu Pro Val Ser Gly Ala Lys Thr Phe Lys Asn Glu Ser
 95 100 105
 Glu Asn Thr Cys Gln Asp Val Asp Glu Cys Gln Gln Asn Pro Arg
 110 115 120
 Leu Cys Lys Ser Tyr Gly Thr Cys Val Asn Thr Leu Gly Ser Tyr
 125 130 135
 Thr Cys Gln Cys Leu Pro Gly Phe Lys Asp Val Asn Glu Cys Thr
 140 145 150
 Ser Gly Gln Asn Pro Cys His Ser Ser Thr His Cys Leu Asn Asn
 155 160 165
 Val Gly Ser Tyr Gln Cys Arg Cys Arg Pro Gly Trp Gln Pro Ile
 170 175 180
 Pro Gly Ser Pro Asn Gly Pro Asn Asn Thr Val Cys Glu Asp Val
 185 190 195
 Asp Glu Cys Ser Ser Gly Gln His Gln Cys Asp Ser Ser Thr Val
 200 205 210
 Cys Phe Asn Thr Val Gly Ser Tyr Ser Cys Arg Cys Arg Pro Gly
 215 220 225
 Trp Lys Pro Arg His Gly Ile Pro Asn Asn Gln Lys Asp Thr Val
 230 235 240
 Cys Glu Asp Met Thr Phe Ser Thr Trp Thr Pro Pro Pro Gly Val

	245		250		255
His Ser Gln Thr	Leu Ser Arg Phe Phe	Asp Lys Val Gln Asp	Leu		
	260		265		270
Gly Arg Asp His	Leu Ser Ser Phe Ala	Val Leu Met Ala His	Tyr		
	275		280		285
Asp Val					



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, 5/10, 1/21, C07K 14/705, 14/72, 16/28, A61K 38/17, C12Q 1/68		A3	(11) International Publication Number: WO 00/20590
			(43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/23317		1240 Dale Avenue #30, Mountain View, CA 94040 (US). GUEGLER, Karl, J. [CH/US]; 1048 Oakland Avenue, Menlo Park, CA 94025 (US). GORGONE, Gina, A. [US/US]; 1253 Pinecrest Drive, Boulder Creek, CA 95006 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 October 1999 (06.10.99)			
(30) Priority Data:			
09/167,219	6 October 1998 (06.10.98) US		
60/172,211	6 October 1998 (06.10.98) US		
60/133,585	11 May 1999 (11.05.99) US		
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications			
US	09/167,219 (CIP)		
Filed on	6 October 1998 (06.10.98)		
US	60/172,211 (CIP)		
Filed on	6 October 1998 (06.10.98)		
US	60/133,585 (CIP)		
Filed on	11 May 1999 (11.05.99)		
(71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). AU-YOUNG, Janice [US/US]; 1419 Kains Avenue, Berkeley, CA 94702 (US). REDDY, Roopa [IN/US]; 1233 W. McKinley Drive, Sunnyvale, CA 94086 (US). CORLEY, Neil, C. [US/US];			

Published

With international search report.

(88) Date of publication of the international search report:

20 July 2000 (20.07.00)

(54) Title: G-PROTEIN COUPLED RECEPTOR PROTEINS

```

1  M-----TPQSLLQTTLFLLSL---LFLV 2763296
1  M-----TPQSLLQTTLFLLSL---LFLV 1595762
1  M V F S V R Q C G H V G R T E E V L L T F K I P L V I I C L H V V L V GI 2117161

21  QGAHGRGRHREDFR-----FCSQRNQTHRSS 2763296
21  QGAHGRGRHREDFR-----FCSQRNQTHRSS 1595762
36  TSLEEDTDNSSLSPPPAKLSVVSFAFPSSN|EVETT|S GI 2117161

45  LHY-----2763296
45  LHYKPTPDLRISIENSEEALTVHAPFPAHPASR| 1595762
71  L N - D V T L S L L P S N E T E K T K I T I V K T P N A S G V K P Q R GI 2117161

49  -----2763296
80  -----SFPDPRGLYHF-----CLYWNRH| 1595762
105 N I C N L S S I C N D S A F F R G E I M F Q Y D K E S T V P Q N Q H I GI 2117161

49  -----2763296
98  -----A G R L H L L Y G K R D -----F L L | 1595762
140 T N G T L T G V L S L S E L K R S E L N K T L Q T L S E T Y F I M C A GI 2117161

49  -----2763296
113 -S D K A S S L L C-----1595762
175 T A E A Q S T L N C T F T I K L N N T M N A C A A I A A L E R V K I R GI 2117161

```

(57) Abstract

The invention provides human G-protein coupled receptor proteins (GCRP) and polynucleotides which identify and encode GCRP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of GCRP.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/US 99/23317

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N5/10 C12N1/21 C07K14/705 C07K14/72
 C07K16/28 A61K38/17 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBEST30 [Online] E.M.B.L. Databases Accession Number: C17798, 4 October 1996 (1996-10-04) FUJIWARA T ET AL: "Human placenta cDNA 5'-end GEN-553A03" XP002130338 99.8% identity in 556 bp overlap with SeqIdNo.9 abstract	3-6
P,X	WO 99 46281 A (BAKER KEVIN P ;CHEN JIAN (US); GENENTECH INC (US); GURNEY AUSTIN () 16 September 1999 (1999-09-16) Human PRO1083: 100.0% identity in 693 aa overlap with SeqIdNo.1 / 99.9% identity in 3574 bp overlap with SeqIdNo.9 --- -/--	1-16,19

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

11 February 2000

Date of mailing of the international search report

16. 05. 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Lonnoy, 0

INTERNATIONAL SEARCH REPORT

Intern: .al Application No

PCT/US 99/23317

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 15551 A (GARVAN INST MED RES ;HERZOG HERBERT (AU)) 1 April 1999 (1999-04-01) TSR32: 99.9% identity in 693 aa overlap with SeqIdNo.1 / 99.7% identity in 2699 bp overlap with SeqIdNo.9 ---	1-16,19
A	EP 0 805 204 A (INST HORMON UND FORTPFLANZUNGS) 5 November 1997 (1997-11-05) ---	
A	WO 96 18651 A (ELLIS CATHERINE ELIZABETH ;SMITHKLINE BEECHAM CORP (US); BERGSMA D) 20 June 1996 (1996-06-20) ---	
A	STADEL J M ET AL: "Orphan G protein-coupled receptors: a neglected opportunity for pioneer drug discovery" TRENDS IN PHARMACOLOGICAL SCIENCES,GB,ELSEVIER TRENDS JOURNAL, CAMBRIDGE, vol. 18, no. 11, November 1997 (1997-11), page 430-437 XP004096215 ISSN: 0165-6147 -----	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/23317

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 19 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 17,18,20
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see additional sheet, invention 1.

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 17,18,20

Claims 17, 18 and 20 refer to an agonist, an antagonist and to the use of an antagonist of the polypeptide of claim 1 without giving a true technical characterization. Moreover, no such compound is defined in the application. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported (Art. 5 and 6 PCT). No search can be carried out for such speculative claims the wording of which is, in fact, a mere recitation of the results to be achieved.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Invention 1: claims 1-20 all partially

A substantially purified polypeptide comprising an amino acid sequence of SeqIdNo.1, a fragment of said polypeptide, a variant polypeptide having at least 90% sequence identity to said polypeptide; an isolated and purified polynucleotide encoding said polypeptide, a variant polynucleotide having at least 70% sequence identity to said polynucleotide, a polynucleotide hybridising to said polynucleotide, a polynucleotide having a sequence complementary to said polynucleotide; a detection method using said complementary polynucleotide; a polynucleotide comprising a sequence of SeqIdNo.9 or a fragment thereof, a polynucleotide having at least 70% sequence identity to said polynucleotide of SeqIdNo.9, a polynucleotide complementary to said polynucleotide of SeqIdNo.9; expression vector, host and recombinant method related thereto; a pharmaceutical composition comprising said polypeptide of SeqIdNo.1 or fragment of said polypeptide of SeqIdNo.1; an antibody to said polypeptide of SeqIdNo.1 or fragment of said polypeptide of SeqIdNo.1; an agonist to said polypeptide of SeqIdNo.1; an antagonist to said polypeptide of SeqIdNo.1; therapeutic application thereof

2. Inventions 2-7: claims 1-20 all partially

Idem as for subject 1 but limited to each of the polypeptide sequences as in SeqIdNo.3-8 and the corresponding polynucleotide sequences as in SeqIdNo.11-16, wherein respectively invention 2 is limited to SeqIdNo.3 and SeqIdNo.11, invention 3 is limited to SeqIdNo.4 and SeqIdNo.12, ..., invention 7 is limited to SeqIdNo.8 and SeqIdNo.16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/23317

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9946281 A	16-09-1999	AU 3072199 A	27-09-1999
		AU 3075099 A	11-10-1999
		WO 9947677 A	23-09-1999
		AU 1532499 A	15-06-1999
		WO 9927098 A	03-06-1999
		AU 3757099 A	08-11-1999
		WO 9954467 A	28-10-1999
		AU 1070399 A	10-05-1999
		WO 9920756 A	29-04-1999

WO 9915551 A	01-04-1999	AU 9330098 A	12-04-1999
		EP 0973800 A	26-01-2000

EP 0805204 A	05-11-1997	DE 19617940 A	30-10-1997

WO 9618651 A	20-06-1996	AU 4522796 A	03-07-1996
		EP 0871669 A	21-10-1998
		JP 10511936 T	17-11-1998
		ZA 9510686 A	03-07-1996
